

Introduction

Modelling parallel systems

Linear Time Properties

Regular Properties

Linear Temporal Logic (LTL)


Computation-Tree Logic

Equivalences and Abstraction



Trace equivalence




BSEQOR5.1-2

$\mathcal{T}_1$:




$\mathcal{T}_2$:




$\mathcal{T}_1$: $\mathcal{T}_2$:

$$\text{Traces}(\mathcal{T}_1) = \{ \emptyset \emptyset a^\omega, \emptyset \emptyset b^\omega \} = \text{Traces}(\mathcal{T}_2)$$



$$\text{Traces}(\mathcal{T}_1) = \{ \emptyset \emptyset a^\omega, \emptyset \emptyset b^\omega \} = \text{Traces}(\mathcal{T}_2)$$

$$\text{CTL-formula } \phi = \exists \text{O}(\exists \text{O} a \wedge \exists \text{O} b)$$




$$\text{Traces}(\mathcal{T}_1) = \{ \emptyset \emptyset a^\omega, \emptyset \emptyset b^\omega \} = \text{Traces}(\mathcal{T}_2)$$

$$\text{CTL-formula } \phi = \exists \text{O}(\exists \text{O} a \wedge \exists \text{O} b)$$

$$\mathcal{T}_1 \not\models \phi \quad \text{and} \quad \mathcal{T}_2 \models \phi$$

Trace equivalence is not compatible with CTL BSEQOR5.1-2



$$\text{Traces}(\mathcal{T}_1) = \{ \emptyset \emptyset a^\omega, \emptyset \emptyset b^\omega \} = \text{Traces}(\mathcal{T}_2)$$

$$\text{CTL-formula } \phi = \exists \text{O}(\exists \text{O} a \wedge \exists \text{O} b)$$

$$\mathcal{T}_1 \not\models \phi \quad \text{and} \quad \mathcal{T}_2 \models \phi$$

- linear vs. branching time
 - * linear time: trace relations
 - * branching time: (bi)simulation relations

- **linear** vs. **branching time**
 - * linear time: trace relations
 - * branching time: (bi)simulation relations
- **(nonsymmetric) preorders** vs. **equivalences**:
 - * preorders: trace inclusion, simulation
 - * equivalences: trace equivalence, bisimulation

- **linear** vs. **branching time**
 - * linear time: trace relations
 - * branching time: (bi)simulation relations
- **(nonsymmetric) preorders** vs. **equivalences**:
 - * preorders: trace inclusion, simulation
 - * equivalences: trace equivalence, bisimulation
- **strong** vs. **weak** relations
 - * strong: reasoning about **all transitions**
 - * weak: abstraction from **stutter steps**

Introduction

Modelling parallel systems

Linear Time Properties

Regular Properties

Linear Temporal Logic (LTL)

Computation-Tree Logic

Equivalences and Abstraction

bisimulation



CTL, CTL*-equivalence

computing the bisimulation quotient

abstraction stutter steps

simulation relations

let $\mathcal{T}_1 = (S_1, Act_1, \rightarrow_1, S_{0,1}, AP, L_1)$,
 $\mathcal{T}_2 = (S_2, Act_2, \rightarrow_2, S_{0,2}, AP, L_2)$

be two transition systems

let $\mathcal{T}_1 = (S_1, Act_1, \rightarrow_1, S_{0,1}, AP, L_1)$,
 $\mathcal{T}_2 = (S_2, Act_2, \rightarrow_2, S_{0,2}, AP, L_2)$

be two transition systems

- with the same set AP

let $\mathcal{T}_1 = (S_1, Act_1, \rightarrow_1, S_{0,1}, AP, L_1)$,
 $\mathcal{T}_2 = (S_2, Act_2, \rightarrow_2, S_{0,2}, AP, L_2)$

be two transition systems

- with the same set AP
- possibly containing terminal states

let $\mathcal{T}_1 = (S_1, Act_1, \rightarrow_1, S_{0,1}, AP, L_1)$,
 $\mathcal{T}_2 = (S_2, Act_2, \rightarrow_2, S_{0,2}, AP, L_2)$

be two transition systems

- with the same set AP
- possibly containing terminal states

Bisimulation equivalence of $\mathcal{T}_1$ and $\mathcal{T}_2$ requires that $\mathcal{T}_1$ and $\mathcal{T}_2$ can simulate each other in a stepwise manner.

$$\text{let } \mathcal{T}_1 = (\mathcal{S}_1, \cancel{\text{Act}_1}, \rightarrow_1, \mathcal{S}_{0,1}, \textcolor{red}{AP}, \mathcal{L}_1), \\ \mathcal{T}_2 = (\mathcal{S}_2, \cancel{\text{Act}_2}, \rightarrow_2, \mathcal{S}_{0,2}, \textcolor{red}{AP}, \mathcal{L}_2)$$

be two transition systems

- with the same set $\textcolor{red}{AP}$ $\longleftarrow$ observables
- possibly containing terminal states

Bisimulation equivalence of $\mathcal{T}_1$ and $\mathcal{T}_2$ requires that $\mathcal{T}_1$ and $\mathcal{T}_2$ can simulate each other in a stepwise manner.

binary relation $\mathcal{R} \subseteq \mathcal{S}_1 \times \mathcal{S}_2$ s.t. for all $(s_1, s_2) \in \mathcal{R}$:

binary relation $\mathcal{R} \subseteq \mathcal{S}_1 \times \mathcal{S}_2$ s.t. for all $(s_1, s_2) \in \mathcal{R}$:

$$(1) \quad L_1(s_1) = L_2(s_2)$$


binary relation $\mathcal{R} \subseteq S_1 \times S_2$ s.t. for all $(s_1, s_2) \in \mathcal{R}$:

$$(1) \quad L_1(s_1) = L_2(s_2)$$


$$(2) \quad \forall s'_1 \in \text{Post}(s_1) \exists s'_2 \in \text{Post}(s_2) \text{ s.t. } (s'_1, s'_2) \in \mathcal{R}$$

binary relation $\mathcal{R} \subseteq S_1 \times S_2$ s.t. for all $(s_1, s_2) \in \mathcal{R}$:

- (1) $L_1(s_1) = L_2(s_2)$
- (2) $\forall s'_1 \in \text{Post}(s_1) \exists s'_2 \in \text{Post}(s_2)$ s.t. $(s'_1, s'_2) \in \mathcal{R}$




can be
completed to




binary relation $\mathcal{R} \subseteq S_1 \times S_2$ s.t. for all $(s_1, s_2) \in \mathcal{R}$:

$$(1) \quad L_1(s_1) = L_2(s_2)$$

$$(2) \quad \forall s'_1 \in \text{Post}(s_1) \exists s'_2 \in \text{Post}(s_2) \text{ s.t. } (s'_1, s'_2) \in \mathcal{R}$$



can be
completed to




$$(3) \quad \forall s'_2 \in \text{Post}(s_2) \exists s'_1 \in \text{Post}(s_1) \text{ s.t. } (s'_1, s'_2) \in \mathcal{R}$$


binary relation $\mathcal{R} \subseteq S_1 \times S_2$ s.t. for all $(s_1, s_2) \in \mathcal{R}$:

$$(1) \quad L_1(s_1) = L_2(s_2)$$

$$(2) \quad \forall s'_1 \in \text{Post}(s_1) \exists s'_2 \in \text{Post}(s_2) \text{ s.t. } (s'_1, s'_2) \in \mathcal{R}$$



can be
completed to



$$(3) \quad \forall s'_2 \in \text{Post}(s_2) \exists s'_1 \in \text{Post}(s_1) \text{ s.t. } (s'_1, s'_2) \in \mathcal{R}$$


and such that the following initial condition holds:

$$(I) \quad \forall s_{0,1} \in S_{0,1} \exists s_{0,2} \in S_{0,2} \text{ s.t. } (s_{0,1}, s_{0,2}) \in \mathcal{R}$$


binary relation $\mathcal{R} \subseteq \mathcal{S}_1 \times \mathcal{S}_2$ s.t. for all $(s_1, s_2) \in \mathcal{R}$:

$$(1) \quad L_1(s_1) = L_2(s_2)$$

$$(2) \quad \forall s'_1 \in \text{Post}(s_1) \exists s'_2 \in \text{Post}(s_2) \text{ s.t. } (s'_1, s'_2) \in \mathcal{R}$$



can be
completed to



$$(3) \quad \forall s'_2 \in \text{Post}(s_2) \exists s'_1 \in \text{Post}(s_1) \text{ s.t. } (s'_1, s'_2) \in \mathcal{R}$$

and such that the following initial condition holds:

$$(I) \quad \forall s_{0,1} \in \mathcal{S}_{0,1} \exists s_{0,2} \in \mathcal{S}_{0,2} \text{ s.t. } (s_{0,1}, s_{0,2}) \in \mathcal{R}$$

$$\forall s_{0,2} \in \mathcal{S}_{0,2} \exists s_{0,1} \in \mathcal{S}_{0,1} \text{ s.t. } (s_{0,1}, s_{0,2}) \in \mathcal{R}$$

bisimulation for $(\mathcal{T}_1, \mathcal{T}_2)$: relation $\mathcal{R} \subseteq S_1 \times S_2$ s.t.

for all $(s_1, s_2) \in \mathcal{R}$:

- (1) labeling condition
- (2) $\left. \begin{array}{l} \text{mutual stepwise} \\ \text{simulation} \end{array} \right\}$
- (3) $\left. \begin{array}{l} \text{mutual stepwise} \\ \text{simulation} \end{array} \right\}$

and initial condition (I)

bisimulation for $(\mathcal{T}_1, \mathcal{T}_2)$: relation $\mathcal{R} \subseteq S_1 \times S_2$ s.t.

for all $(s_1, s_2) \in \mathcal{R}$:

- (1) labeling condition
- (2) $\left. \begin{array}{l} \text{mutual stepwise} \\ \text{simulation} \end{array} \right\}$
- (3)

and initial condition (I)

bisimulation equivalence $\sim$ for TS:

bisimulation for $(\mathcal{T}_1, \mathcal{T}_2)$: relation $\mathcal{R} \subseteq S_1 \times S_2$ s.t.

for all $(s_1, s_2) \in \mathcal{R}$:

- (1) labeling condition
- (2) $\left. \begin{array}{l} \text{mutual stepwise} \\ \text{simulation} \end{array} \right\}$
- (3)

and initial condition (I)

bisimulation equivalence $\sim$ for TS:

$\mathcal{T}_1 \sim \mathcal{T}_2$ iff there is a bisimulation $\mathcal{R}$ for $(\mathcal{T}_1, \mathcal{T}_2)$

bisimulation for $(\mathcal{T}_1, \mathcal{T}_2)$: relation $\mathcal{R} \subseteq S_1 \times S_2$ s.t.

for all $(s_1, s_2) \in \mathcal{R}$:

- (1) labeling condition
- (2) $\left. \begin{array}{l} \text{ } \end{array} \right\}$ mutual stepwise
- (3) $\left. \begin{array}{l} \text{ } \end{array} \right\}$ simulation

and initial condition (I)

bisimulation equivalence $\sim$ for TS:



$\mathcal{T}_1 \sim \mathcal{T}_2$ iff there is a bisimulation $\mathcal{R}$ for $(\mathcal{T}_1, \mathcal{T}_2)$

for state s_1 of $\mathcal{T}_1$ and state s_2 of $\mathcal{T}_2$:

$s_1 \sim s_2$ iff there exists a bisimulation $\mathcal{R}$ for $(\mathcal{T}_1, \mathcal{T}_2)$
such that $(s_1, s_2) \in \mathcal{R}$

Two beverage machines



BSEQOR5.1-8-BIS



$$AP = \{ \text{pay}, \text{coke}, \text{soda} \}$$

Two beverage machines



BSEQOR5.1-8-BIS



$$AP = \{ \textit{pay}, \textit{coke}, \textit{soda} \}$$

Two beverage machines

BSEQOR5.1-8-BIS




$$AP = \{ \textit{pay}, \textit{coke}, \textit{soda} \}$$

$$\mathcal{T}_1 \sim \mathcal{T}_2$$

Two beverage machines

BSEQOR5.1-8-BIS




$$AP = \{ \text{pay}, \text{coke}, \text{soda} \}$$

$\mathcal{T}_1 \sim \mathcal{T}_2$ as there is a bisimulation for $(\mathcal{T}_1, \mathcal{T}_2)$:

Two beverage machines

BSEQOR5.1-8-BIS





$$AP = \{ \text{pay}, \text{coke}, \text{soda} \}$$

$\mathcal{T}_1 \sim \mathcal{T}_2$ as there is a bisimulation for $(\mathcal{T}_1, \mathcal{T}_2)$:

$$\left\{ \begin{array}{l} (\text{pay}, \text{pay}), (\text{select}, \text{select}), (\text{soda}, \text{soda}) \\ (\text{coke}, \text{coke}_1), (\text{coke}, \text{coke}_2) \end{array} \right\}$$

Two beverage machines



BSEQOR5.1-8-BIS-3



$$AP = \{\textit{pay}, \textit{coke}, \textit{soda}\}$$

Two beverage machines

BSEQOR5.1-8-BIS-3





$AP = \{\text{pay}, \text{coke}, \text{soda}\}$

$\mathcal{T}_1 \not\sim \mathcal{T}_2$

Two beverage machines

BSEQOR5.1-8-BIS-3




$AP = \{\text{pay}, \text{coke}, \text{soda}\}$


$\mathcal{T}_1 \not\sim \mathcal{T}_2$


because there is no state in $\mathcal{T}_1$ that has both


- a successor labeled with **coke** and
- a successor labeled with **soda**



can be
completed to









can be
completed to

Path lifting for bisimulation $\mathcal{R}$

BSEQOR5.1-9-BIS




can be
completed to




Path lifting for bisimulation $\mathcal{R}$

BSEQOR5.1-9-BIS




can be
completed to




Path lifting for bisimulation $\mathcal{R}$

BSEQOR5.1-9-BIS




can be
completed to




Path lifting for bisimulation $\mathcal{R}$

BSEQOR5.1-9-BIS



can be
completed to



Properties of bisimulation equivalence

BSEQOR5.1-PROP-OF-BIS.TEX

$\sim$ is an **equivalence**

$\sim$ is an **equivalence**, i.e.,

- reflexivity: $\mathcal{T} \sim \mathcal{T}$ for all transition systems $\mathcal{T}$

$\sim$ is an **equivalence**, i.e.,

- reflexivity: $\mathcal{T} \sim \mathcal{T}$ for all transition systems $\mathcal{T}$



If S is the state space of $\mathcal{T}$ then

$$\mathcal{R} = \{(s, s) : s \in S\}$$

is a bisimulation for $(\mathcal{T}, \mathcal{T})$

$\sim$ is an **equivalence**, i.e.,

- reflexivity: $\mathcal{T} \sim \mathcal{T}$ for all transition systems $\mathcal{T}$
- symmetry: $\mathcal{T}_1 \sim \mathcal{T}_2$ implies $\mathcal{T}_2 \sim \mathcal{T}_1$

$\sim$ is an **equivalence**, i.e.,

- reflexivity: $\mathcal{T} \sim \mathcal{T}$ for all transition systems $\mathcal{T}$
- symmetry: $\mathcal{T}_1 \sim \mathcal{T}_2$ implies $\mathcal{T}_2 \sim \mathcal{T}_1$



If $\mathcal{R}$ is a bisimulation for $(\mathcal{T}_1, \mathcal{T}_2)$ then

$$\mathcal{R}^{-1} = \{(s_2, s_1) : (s_1, s_2) \in \mathcal{R}\}$$

is a bisimulation for $(\mathcal{T}_2, \mathcal{T}_1)$

$\sim$ is an **equivalence**, i.e.,

- reflexivity: $\mathcal{T} \sim \mathcal{T}$ for all transition systems $\mathcal{T}$
- symmetry: $\mathcal{T}_1 \sim \mathcal{T}_2$ implies $\mathcal{T}_2 \sim \mathcal{T}_1$
- transitivity: if $\mathcal{T}_1 \sim \mathcal{T}_2$ and $\mathcal{T}_2 \sim \mathcal{T}_3$ then $\mathcal{T}_1 \sim \mathcal{T}_3$

$\sim$ is an **equivalence**, i.e.,

- reflexivity: $\mathcal{T} \sim \mathcal{T}$ for all transition systems $\mathcal{T}$
- symmetry: $\mathcal{T}_1 \sim \mathcal{T}_2$ implies $\mathcal{T}_2 \sim \mathcal{T}_1$
- transitivity: if $\mathcal{T}_1 \sim \mathcal{T}_2$ and $\mathcal{T}_2 \sim \mathcal{T}_3$ then $\mathcal{T}_1 \sim \mathcal{T}_3$



Let $\mathcal{R}_{1,2}$ be a bisimulation for $(\mathcal{T}_1, \mathcal{T}_2)$,
 $\mathcal{R}_{2,3}$ be a bisimulation for $(\mathcal{T}_2, \mathcal{T}_3)$.

$\sim$ is an **equivalence**, i.e.,

- reflexivity: $\mathcal{T} \sim \mathcal{T}$ for all transition systems $\mathcal{T}$
- symmetry: $\mathcal{T}_1 \sim \mathcal{T}_2$ implies $\mathcal{T}_2 \sim \mathcal{T}_1$
- transitivity: if $\mathcal{T}_1 \sim \mathcal{T}_2$ and $\mathcal{T}_2 \sim \mathcal{T}_3$ then $\mathcal{T}_1 \sim \mathcal{T}_3$



Let $\mathcal{R}_{1,2}$ be a bisimulation for $(\mathcal{T}_1, \mathcal{T}_2)$,


$\mathcal{R}_{2,3}$ be a bisimulation for $(\mathcal{T}_2, \mathcal{T}_3)$.

$$\mathcal{R} \stackrel{\text{def}}{=} \left\{ (s_1, s_3) : \exists s_2 \text{ s.t. } (s_1, s_2) \in \mathcal{R}_{1,2} \right. \\ \left. \text{and } (s_2, s_3) \in \mathcal{R}_{2,3} \right\}$$


is a bisimulation for $(\mathcal{T}_1, \mathcal{T}_3)$

Correct or wrong?

BSEQOR5.1-19




~




Correct or wrong?

BSEQOR5.1-19




$\sim$




wrong

Correct or wrong?

BSEQOR5.1-19



$\sim$




wrong


$s_1 \longrightarrow u$, but $s_2 \not\longrightarrow \text{blue}$ (thus $s_1 \not\sim s_2$)

Correct or wrong?

BSEQOR5.1-19




$\sim$




wrong

$s_1 \longrightarrow u$, but $s_2 \not\longrightarrow \text{blue}$ (thus $s_1 \not\sim s_2$)




$\sim$




Correct or wrong?

BSEQOR5.1-19




$\sim$




wrong

$s_1 \rightarrow u$, but $s_2 \not\rightarrow \text{blue}$ (thus $s_1 \not\sim s_2$)




$\sim$




correct

Correct or wrong?

BSEQOR5.1-19




$\sim$




wrong

$s_1 \longrightarrow u$, but $s_2 \not\rightarrow \text{blue}$ (thus $s_1 \not\sim s_2$)



$\sim$




correct

bisimulation:

$\{(w_1, w_2), (w_1', w_2), (s_1, s_2'), (s_1, s_2'), (u, x), (u, y)\}$


Correct or wrong?

BSEQOR5.1-20




Correct or wrong?

BSEQOR5.1-20




~




correct

Correct or wrong?

BSEQOR5.1-20



$\sim$




correct

bisimulation


$$\{(s_1, s_2), (s_1', s_2'), (s_1', s_2''), (t_1, t_2), (t_1', t_2), (t_1'', t_2)\}$$

Correct or wrong?

BSEQOR5.1-20




$\sim$




correct

bisimulation

$$\{(s_1, s_2), (s'_1, s'_2), (s'_1, s''_2), (t_1, t_2), (t'_1, t_2), (t''_1, t_2)\}$$




$\sim$




Correct or wrong?

BSEQOR5.1-20




$\sim$




correct

bisimulation

$$\{(s_1, s_2), (s'_1, s'_2), (s'_1, s''_2), (t_1, t_2), (t'_1, t_2), (t''_1, t_2)\}$$





$\sim$



correct

Correct or wrong?

BSEQOR5.1-20



bisimulation: $\{(s_1, s_2), (t_1, t_2), (t'_1, t_2), (u_1, u_2), (v_1, v_2)\}$

$$\mathcal{T}_1 \sim \mathcal{T}_2 \implies \textit{Traces}(\mathcal{T}_1) = \textit{Traces}(\mathcal{T}_2)$$

$$\mathcal{T}_1 \sim \mathcal{T}_2 \implies \text{Traces}(\mathcal{T}_1) = \text{Traces}(\mathcal{T}_2)$$

proof: ... path fragment lifting ...

$$\mathcal{T}_1 \sim \mathcal{T}_2 \implies \text{Traces}(\mathcal{T}_1) = \text{Traces}(\mathcal{T}_2)$$


proof: ... path fragment lifting ...

$$\text{Traces}(\mathcal{T}_1) = \text{Traces}(\mathcal{T}_2) \not\Rightarrow \mathcal{T}_1 \sim \mathcal{T}_2$$

$$\mathcal{T}_1 \sim \mathcal{T}_2 \implies \text{Traces}(\mathcal{T}_1) = \text{Traces}(\mathcal{T}_2)$$

proof: ... path fragment lifting ...

$$\text{Traces}(\mathcal{T}_1) = \text{Traces}(\mathcal{T}_2) \not\Rightarrow \mathcal{T}_1 \sim \mathcal{T}_2$$



trace equivalent, but not bisimulation equivalent

$$\mathcal{T}_1 \sim \mathcal{T}_2 \implies \text{Traces}(\mathcal{T}_1) = \text{Traces}(\mathcal{T}_2)$$

proof: ... path fragment lifting ...

$$\text{Traces}(\mathcal{T}_1) = \text{Traces}(\mathcal{T}_2) \not\Rightarrow \mathcal{T}_1 \sim \mathcal{T}_2$$

Trace equivalence is **strictly coarser** than
bisimulation equivalence.

$$\mathcal{T}_1 \sim \mathcal{T}_2 \implies \text{Traces}(\mathcal{T}_1) = \text{Traces}(\mathcal{T}_2)$$

proof: ... path fragment lifting ...



$$\text{Traces}(\mathcal{T}_1) = \text{Traces}(\mathcal{T}_2) \not\Rightarrow \mathcal{T}_1 \sim \mathcal{T}_2$$

Trace equivalence is **strictly coarser** than
bisimulation equivalence.



Bisimulation equivalent transition systems satisfy
the **same LT properties** (e.g., **LTL formulas**).

- as a relation that compares **2** transition systems

- as a relation that compares **2** transition systems



 $\mathcal{T}_1$  $\mathcal{T}_2$ 

- as a relation that compares **2** transition systems


 $\mathcal{T}_1$  $\mathcal{T}_2$ 

- as a relation on the **states** of **1** transition system

- as a relation that compares **2** transition systems




- as a relation on the **states** of **1** transition system



- as a relation that compares **2** transition systems



- as a relation on the **states** of **1** transition system




$$s_1 \sim s_2 \text{ iff } \mathcal{T}_{s_1} \sim \mathcal{T}_{s_2}$$

- as a relation that compares **2** transition systems



- as a relation on the **states** of **1** transition system




$$s_1 \sim s_2 \text{ iff } \mathcal{T}_{s_1} \sim \mathcal{T}_{s_2}$$

- as a relation that compares **2** transition systems



- as a relation on the **states** of **1** transition system



$s_1 \sim s_2$ iff $\mathcal{T}_{s_1} \sim \mathcal{T}_{s_2}$ iff
there exists a bisimulation $\mathcal{R}$ for $\mathcal{T}$ s.t. $(s_1, s_2) \in \mathcal{R}$

Let $\mathcal{T}$ be a TS with proposition set AP .

Let $\mathcal{T}$ be a TS with proposition set AP .

A **bisimulation** for $\mathcal{T}$ is a binary relation $\mathcal{R}$ on the state space of $\mathcal{T}$ s.t. for all $(s_1, s_2) \in \mathcal{R}$:

- (1) $L(s_1) = L(s_2)$
- (2) $\forall s'_1 \in \text{Post}(s_1) \exists s'_2 \in \text{Post}(s_2) \text{ s.t. } (s'_1, s'_2) \in \mathcal{R}$
- (3) $\forall s'_2 \in \text{Post}(s_2) \exists s'_1 \in \text{Post}(s_1) \text{ s.t. } (s'_1, s'_2) \in \mathcal{R}$

Let $\mathcal{T}$ be a TS with proposition set AP .

A **bisimulation** for $\mathcal{T}$ is a binary relation $\mathcal{R}$ on the state space of $\mathcal{T}$ s.t. for all $(s_1, s_2) \in \mathcal{R}$:

- (1) $L(s_1) = L(s_2)$
- (2) $\forall s'_1 \in \text{Post}(s_1) \exists s'_2 \in \text{Post}(s_2)$ s.t. $(s'_1, s'_2) \in \mathcal{R}$
- (3) $\forall s'_2 \in \text{Post}(s_2) \exists s'_1 \in \text{Post}(s_1)$ s.t. $(s'_1, s'_2) \in \mathcal{R}$

bisimulation equivalence $\sim_{\mathcal{T}}$:

$s_1 \sim_{\mathcal{T}} s_2$ iff there exists a bisimulation $\mathcal{R}$ for $\mathcal{T}$
s.t. $(s_1, s_2) \in \mathcal{R}$

Let $\mathcal{T}$ be a TS with proposition set AP .

A **bisimulation** for $\mathcal{T}$ is a binary relation $\mathcal{R}$ on the state space of $\mathcal{T}$ s.t. for all $(s_1, s_2) \in \mathcal{R}$:

- (1) $L(s_1) = L(s_2)$
- (2) $\forall s'_1 \in \text{Post}(s_1) \exists s'_2 \in \text{Post}(s_2)$ s.t. $(s'_1, s'_2) \in \mathcal{R}$
- (3) $\forall s'_2 \in \text{Post}(s_2) \exists s'_1 \in \text{Post}(s_1)$ s.t. $(s'_1, s'_2) \in \mathcal{R}$

coinductive definition of $\sim_{\mathcal{T}}$:

$s_1 \sim_{\mathcal{T}} s_2$ iff there exists a bisimulation $\mathcal{R}$ for $\mathcal{T}$
s.t. $(s_1, s_2) \in \mathcal{R}$

Let $\mathcal{T}$ be a transition system with state space S .

Bisimulation equivalence $\sim_{\mathcal{T}}$ is

Let $\mathcal{T}$ be a transition system with state space S .

Bisimulation equivalence $\sim_{\mathcal{T}}$ is

- the coarsest bisimulation on $\mathcal{T}$

Let $\mathcal{T}$ be a transition system with state space S .

Bisimulation equivalence $\sim_{\mathcal{T}}$ is

- the coarsest bisimulation on $\mathcal{T}$
- and an equivalence on S

Let $\mathcal{T}$ be a transition system with state space S .

Bisimulation equivalence $\sim_{\mathcal{T}}$ is the coarsest equivalence on S s.t. for all states $s_1, s_2 \in S$ with $s_1 \sim_{\mathcal{T}} s_2$:

Let $\mathcal{T}$ be a transition system with state space S .


Bisimulation equivalence $\sim_{\mathcal{T}}$ is the coarsest equivalence on S s.t. for all states $s_1, s_2 \in S$ with $s_1 \sim_{\mathcal{T}} s_2$:

- (1) $L(s_1) = L(s_2)$
- (2) each transition of s_1 can be mimicked by a transition of s_2 :


Let $\mathcal{T}$ be a transition system with state space $\mathcal{S}$.

Bisimulation equivalence $\sim_{\mathcal{T}}$ is the **coarsest equivalence** on $\mathcal{S}$ s.t. for all states $s_1, s_2 \in \mathcal{S}$ with $s_1 \sim_{\mathcal{T}} s_2$:

- (1) $L(s_1) = L(s_2)$
- (2) each transition of s_1 can be mimicked by a transition of s_2 :



can be
completed to



Two variants of bisimulation equivalence

BSEQOR5.1-31

- $\sim$ relation that compares **2** transition systems
- $\sim_{\mathcal{T}}$ equivalence on the state space of a single TS $\mathcal{T}$

Two variants of bisimulation equivalence

BSEQOR5.1-31

- $\sim$ relation that compares **2** transition systems
 - $\sim_{\mathcal{T}}$ equivalence on the state space of a single TS $\mathcal{T}$
1. $\sim_{\mathcal{T}}$ can be derived from $\sim$

Two variants of bisimulation equivalence

BSEQOR5.1-31

- $\sim$ relation that compares **2** transition systems
- $\sim_{\mathcal{T}}$ equivalence on the state space of a single TS $\mathcal{T}$

1. $\sim_{\mathcal{T}}$ can be derived from $\sim$

for all states s_1 and s_2 of $\mathcal{T}$:

$$s_1 \sim_{\mathcal{T}} s_2 \quad \text{iff} \quad \mathcal{T}_{s_1} \sim \mathcal{T}_{s_2}$$

Two variants of bisimulation equivalence

BSEQOR5.1-31

- $\sim$ relation that compares **2** transition systems
- $\sim_{\mathcal{T}}$ equivalence on the state space of a single TS $\mathcal{T}$

1. $\sim_{\mathcal{T}}$ can be derived from $\sim$

for all states s_1 and s_2 of $\mathcal{T}$:

$$s_1 \sim_{\mathcal{T}} s_2 \quad \text{iff} \quad \mathcal{T}_{s_1} \sim \mathcal{T}_{s_2}$$



where $\mathcal{T}_s$ agrees with $\mathcal{T}$, except that state s is declared to be the unique initial state

Two variants of bisimulation equivalence

BSEQOR5.1-31

- $\sim$ relation that compares **2** transition systems
- $\sim_{\mathcal{T}}$ equivalence on the state space of a single TS $\mathcal{T}$

1. $\sim_{\mathcal{T}}$ can be derived from $\sim$

for all states s_1 and s_2 of $\mathcal{T}$:

$$s_1 \sim_{\mathcal{T}} s_2 \quad \text{iff} \quad \mathcal{T}_{s_1} \sim \mathcal{T}_{s_2}$$




where $\mathcal{T}_s$ agrees with $\mathcal{T}$, except that state s is declared to be the unique initial state

2. $\sim$ can be derived from $\sim_{\mathcal{T}}$

given two transition systems $\mathcal{T}_1$ and $\mathcal{T}_2$

$\mathcal{T}_1$ with state space S_1




$\mathcal{T}_2$ with state space S_2




given two transition systems $\mathcal{T}_1$ and $\mathcal{T}_2$

$\mathcal{T}_1$ with state space S_1




$\mathcal{T}_2$ with state space S_2




consider $\mathcal{T} = \mathcal{T}_1 \uplus \mathcal{T}_2$
(state space $S_1 \uplus S_2$)

given two transition systems $\mathcal{T}_1$ and $\mathcal{T}_2$

$\mathcal{T}_1$ with state space S_1



$\mathcal{T}_2$ with state space S_2




consider $\mathcal{T} = \mathcal{T}_1 \uplus \mathcal{T}_2$
(state space $S_1 \uplus S_2$)

$\mathcal{T}_1 \sim \mathcal{T}_2$ iff $\forall$ initial states s_1 of $\mathcal{T}_1$
 $\exists$ initial state s_2 of $\mathcal{T}_2$ s.t. $s_1 \sim_{\mathcal{T}} s_2$,

given two transition systems $\mathcal{T}_1$ and $\mathcal{T}_2$


$\mathcal{T}_1$ with state space S_1



$\mathcal{T}_2$ with state space S_2



consider $\mathcal{T} = \mathcal{T}_1 \uplus \mathcal{T}_2$
(state space $S_1 \uplus S_2$)



$\mathcal{T}_1 \sim \mathcal{T}_2$ iff $\forall$ initial states s_1 of $\mathcal{T}_1$
 $\exists$ initial state s_2 of $\mathcal{T}_2$ s.t. $s_1 \sim_{\mathcal{T}} s_2$,
and vice versa

Let $\mathcal{T} = (\mathcal{S}, \text{Act}, \rightarrow, \mathcal{S}_0, \text{AP}, \mathcal{L})$ be a TS.

Let $\mathcal{T} = (\mathcal{S}, \text{Act}, \rightarrow, \mathcal{S}_0, \text{AP}, \mathcal{L})$ be a TS.

bisimulation quotient $\mathcal{T}/\sim$ arises from $\mathcal{T}$
by collapsing bisimulation equivalent states

Let $\mathcal{T} = (\mathcal{S}, Act, \rightarrow, \mathcal{S}_0, AP, L)$ be a TS.

bisimulation quotient:

$$\mathcal{T}/\sim = (\mathcal{S}', Act', \rightarrow', \mathcal{S}'_0, AP, L')$$

Let $\mathcal{T} = (\mathcal{S}, \text{Act}, \rightarrow, \mathcal{S}_0, \text{AP}, L)$ be a TS.

bisimulation quotient:

$$\mathcal{T}/\sim = (\mathcal{S}', \text{Act}', \rightarrow', \mathcal{S}'_0, \text{AP}, L')$$

- state space: $\mathcal{S}' = \mathcal{S}/\sim_{\mathcal{T}}$



set of bisimulation equivalence classes

Let $\mathcal{T} = (\mathcal{S}, \text{Act}, \rightarrow, \mathcal{S}_0, \text{AP}, L)$ be a TS.

bisimulation quotient:

$$\mathcal{T}/\sim = (\mathcal{S}', \text{Act}', \rightarrow', \mathcal{S}'_0, \text{AP}, L')$$

- state space: $\mathcal{S}' = \mathcal{S}/\sim_{\mathcal{T}}$
- set of initial states: $\mathcal{S}'_0 = \{[s]_{\sim_{\mathcal{T}}} : s \in \mathcal{S}_0\}$

Let $\mathcal{T} = (\mathcal{S}, \text{Act}, \rightarrow, \mathcal{S}_0, \text{AP}, L)$ be a TS.

bisimulation quotient:

$$\mathcal{T}/\sim = (\mathcal{S}', \text{Act}', \rightarrow', \mathcal{S}'_0, \text{AP}, L')$$

- state space: $\mathcal{S}' = \mathcal{S}/\sim_{\mathcal{T}}$
- set of initial states: $\mathcal{S}'_0 = \{[s]_{\sim_{\mathcal{T}}} : s \in \mathcal{S}_0\}$
- labeling function: $L'([s]_{\sim_{\mathcal{T}}}) = L(s)$

Let $\mathcal{T} = (\mathcal{S}, \text{Act}, \rightarrow, \mathcal{S}_0, \text{AP}, L)$ be a TS.

bisimulation quotient:

$$\mathcal{T}/\sim = (\mathcal{S}', \text{Act}', \rightarrow', \mathcal{S}'_0, \text{AP}, L')$$

- state space: $\mathcal{S}' = \mathcal{S}/\sim_{\mathcal{T}}$
- set of initial states: $\mathcal{S}'_0 = \{[s]_{\sim_{\mathcal{T}}} : s \in \mathcal{S}_0\}$
- labeling function: $L'([s]_{\sim_{\mathcal{T}}}) = L(s)$

well-defined

by the labeling condition
of bisimulations

Let $\mathcal{T} = (\mathcal{S}, \text{Act}, \rightarrow, \mathcal{S}_0, \text{AP}, L)$ be a TS.

bisimulation quotient:

$$\mathcal{T}/\sim = (\mathcal{S}', \text{Act}', \rightarrow', \mathcal{S}'_0, \text{AP}, L')$$

- state space: $\mathcal{S}' = \mathcal{S}/\sim_{\mathcal{T}}$
- set of initial states: $\mathcal{S}'_0 = \{[s]_{\sim_{\mathcal{T}}} : s \in \mathcal{S}_0\}$
- labeling function: $L'([s]_{\sim_{\mathcal{T}}}) = L(s)$
- transition relation:

$$\frac{s \longrightarrow s'}{[s]_{\sim_{\mathcal{T}}} \longrightarrow [s']_{\sim_{\mathcal{T}}}}$$

Let $\mathcal{T} = (\mathcal{S}, \text{Act}, \rightarrow, \mathcal{S}_0, \text{AP}, L)$ be a TS.

bisimulation quotient:

$$\mathcal{T}/\sim = (\mathcal{S}', \text{Act}', \rightarrow', \mathcal{S}'_0, \text{AP}, L')$$

- state space: $\mathcal{S}' = \mathcal{S}/\sim_{\mathcal{T}}$
- set of initial states: $\mathcal{S}'_0 = \{[s]_{\sim_{\mathcal{T}}} : s \in \mathcal{S}_0\}$
- labeling function: $L'([s]_{\sim_{\mathcal{T}}}) = L(s)$
- transition relation:

$$\frac{s \longrightarrow s'}{[s]_{\sim_{\mathcal{T}}} \longrightarrow [s']_{\sim_{\mathcal{T}}}}$$

action labels
irrelevant

Let $\mathcal{T} = (\mathcal{S}, \text{Act}, \rightarrow, \mathcal{S}_0, \text{AP}, L)$ be a TS.

bisimulation quotient:

$$\mathcal{T}/\sim = (\mathcal{S}', \{\tau\}, \rightarrow', \mathcal{S}'_0, \text{AP}, L')$$

- state space: $\mathcal{S}' = \mathcal{S}/\sim_{\mathcal{T}}$
- set of initial states: $\mathcal{S}'_0 = \{[s]_{\sim_{\mathcal{T}}} : s \in \mathcal{S}_0\}$
- labeling function: $L'([s]_{\sim_{\mathcal{T}}}) = L(s)$
- transition relation:

$$\frac{s \xrightarrow{\alpha} s'}{[s]_{\sim_{\mathcal{T}}} \xrightarrow{\tau} [s']_{\sim_{\mathcal{T}}}}$$

action labels
irrelevant

Let $\mathcal{T} = (\mathcal{S}, \text{Act}, \rightarrow, \mathcal{S}_0, \text{AP}, \mathcal{L})$ be a TS.

bisimulation quotient:

$$\mathcal{T}/\sim = (\mathcal{S}', \{\tau\}, \rightarrow', \mathcal{S}'_0, \text{AP}, \mathcal{L}')$$

- state space: $\mathcal{S}' = \mathcal{S}/\sim_{\mathcal{T}}$
- set of initial states: $\mathcal{S}'_0 = \{[s]_{\sim_{\mathcal{T}}} : s \in \mathcal{S}_0\}$
- labeling function: $\mathcal{L}'([s]_{\sim_{\mathcal{T}}}) = \mathcal{L}(s)$
- transition relation:

$$\frac{s \xrightarrow{\alpha} s'}{[s]_{\sim_{\mathcal{T}}} \xrightarrow{\tau} [s']_{\sim_{\mathcal{T}}}}$$

$$\mathcal{T} \sim \mathcal{T}/\sim$$

Example: interleaving of n printers

BSEQOR5.1-34


parallel system $\mathcal{T} = \underbrace{\textit{Printer} ||| \textit{Printer} ||| \dots ||| \textit{Printer}}_{n \text{ printers}}$

Example: interleaving of n printers

BSEQOR5.1-34

parallel system $\mathcal{T} = \underbrace{\textit{Printer} ||| \textit{Printer} ||| \dots ||| \textit{Printer}}_{n \text{ printers}}$

transition system
for each printer




Example: interleaving of n printers

BSEQOR5.1-34

parallel system $\mathcal{T} = \underbrace{\textit{Printer} ||| \textit{Printer} ||| \dots ||| \textit{Printer}}_{n \text{ printers}}$

$AP = \{0, 1, \dots, n\}$ “number of available printers”

transition system
for each printer




Example: $n=3$ printers

BSEQOR5.1-34

parallel system $\mathcal{T} = \underbrace{\textit{Printer} ||| \textit{Printer} ||| \dots ||| \textit{Printer}}_{n \text{ printers}}$

$AP = \{0, 1, 2, 3\}$



p : is printing


r : ready to print

Example: $n=3$ printers

BSEQOR5.1-34

parallel system $\mathcal{T} = \underbrace{\textit{Printer} ||| \textit{Printer} ||| \dots ||| \textit{Printer}}_{n \text{ printers}}$

$AP = \{0, 1, 2, 3\}$



p : is printing


r : ready to print

Example: $n=3$ printers


BSEQOR5.1-34

parallel system $\mathcal{T} = \underbrace{\text{Printer} ||| \text{Printer} ||| \dots ||| \text{Printer}}_{n \text{ printers}}$

$$AP = \{0, 1, 2, 3\}$$



p : is printing
 r : ready to print




bisimulation
quotient

Example: $n=3$ printers


BSEQOR5.1-34

parallel system $\mathcal{T} = \underbrace{\text{Printer} ||| \text{Printer} ||| \dots ||| \text{Printer}}_{n \text{ printers}}$

$$AP = \{0, 1, 2, 3\}$$



2^n states



$n+1$ states

solutions for mutual exclusion problems:

- semaphore
- Peterson's algorithm

solutions for mutual exclusion problems:

- semaphore
- Peterson's algorithm
- Bakery algorithm

solutions for mutual exclusion problems:

- semaphore
- Peterson's algorithm
- Bakery algorithm



given two concurrent processes P_1 and P_2

solutions for mutual exclusion problems:

- semaphore
- Peterson's algorithm
- Bakery algorithm



given two concurrent processes P_1 and P_2

- two additional shared variables: $x_1, x_2 \in \mathbb{N}$

solutions for mutual exclusion problems:

- semaphore
- Peterson's algorithm
- Bakery algorithm



given two concurrent processes P_1 and P_2

- two additional shared variables: $x_1, x_2 \in \mathbb{N}$
- if P_1 and P_2 are waiting then:

solutions for mutual exclusion problems:

- semaphore
- Peterson's algorithm
- Bakery algorithm



given two concurrent processes P_1 and P_2

- two additional shared variables: $x_1, x_2 \in \mathbb{N}$
- if P_1 and P_2 are waiting then:
 - if $x_1 < x_2$ then P_1 enters its critical section
 - if $x_2 < x_1$ then P_2 enters its critical section

solutions for mutual exclusion problems:

- semaphore
- Peterson's algorithm
- Bakery algorithm



given two concurrent processes P_1 and P_2

- two additional shared variables: $x_1, x_2 \in \mathbb{N}$
- if P_1 and P_2 are waiting then:
 - if $x_1 < x_2$ then P_1 enters its critical section
 - if $x_2 < x_1$ then P_2 enters its critical section
 - $x_1 = x_2$: cannot happen

protocol for P_1 :

```
LOOP FOREVER
```

```
  noncritical actions
```

```
   $x_1 := x_2 + 1$ 
```

```
  AWAIT ( $x_1 < x_2$ )  $\vee$  ( $x_2 = 0$ );
```

```
  critical section;
```

```
   $x_1 := 0$ 
```

```
END LOOP
```

symmetric protocol for P_2

protocol for P_1 :

LOOP FOREVER

noncritical actions

$x_1 := x_2 + 1$

AWAIT ($x_1 < x_2$) $\vee$ ($x_2 = 0$);

critical section;

$x_1 := 0$

END LOOP



initially:

$x_1 = x_2 = 0$

symmetric protocol for P_2



Program graphs for the Bakery algorithm

BSEQOR5.1-37





Transition system for the Bakery algorithm

BSEQOR5.1-37





Transition system for the Bakery algorithm

BSEQOR5.1-37




Transition system for the Bakery algorithm

BSEQOR5.1-37





Transition system for the Bakery algorithm

BSEQOR5.1-37





Transition system for the Bakery algorithm



BSEQOR5.1-37



Transition system for the Bakery algorithm

BSEQOR5.1-37






infinite transition system with a
finite bisimulation quotient


Bakery algorithm: bisimulation quotient

BSEQOR5.1-38




Bakery algorithm: bisimulation quotient

BSEQOR5.1-38




Bakery algorithm: bisimulation quotient

BSEQOR5.1-38



Bakery algorithm: bisimulation quotient

BSEQOR5.1-38



Introduction

Modelling parallel systems

Linear Time Properties

Regular Properties

Linear Temporal Logic (LTL)

Computation-Tree Logic

Equivalences and Abstraction

bisimulation

CTL, CTL*-equivalence



computing the bisimulation quotient

abstraction stutter steps

simulation relations

CTL* state formulas

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \exists \varphi$$

CTL* path formulas

$$\varphi ::= \Phi \mid \varphi_1 \wedge \varphi_2 \mid \neg \varphi \mid \bigcirc \varphi \mid \varphi_1 \mathbf{U} \varphi_2$$

CTL* state formulas

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \exists \varphi$$

CTL* path formulas

$$\varphi ::= \Phi \mid \varphi_1 \wedge \varphi_2 \mid \neg \varphi \mid \bigcirc \varphi \mid \varphi_1 \mathbf{U} \varphi_2$$

derived operators:

- $\Diamond, \Box, \dots$ as in **LTL**

CTL* state formulas

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \exists \varphi$$

CTL* path formulas

$$\varphi ::= \Phi \mid \varphi_1 \wedge \varphi_2 \mid \neg \varphi \mid \bigcirc \varphi \mid \varphi_1 \mathbf{U} \varphi_2$$

derived operators:

- $\Diamond, \Box, \dots$ as in **LTL**
- universal quantification: $\forall \varphi \stackrel{\text{def}}{=} \neg \exists \neg \varphi$

CTL* state formulas

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \exists \varphi$$

CTL* path formulas

$$\varphi ::= \Phi \mid \varphi_1 \wedge \varphi_2 \mid \neg \varphi \mid \bigcirc \varphi \mid \varphi_1 \mathbf{U} \varphi_2$$

CTL: sublogic of **CTL***

CTL* state formulas

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \exists \varphi$$

CTL* path formulas

$$\varphi ::= \Phi \mid \varphi_1 \wedge \varphi_2 \mid \neg \varphi \mid \bigcirc \varphi \mid \varphi_1 \mathbf{U} \varphi_2$$

CTL: sublogic of **CTL***

- with path quantifiers $\exists$ and $\forall$

CTL* state formulas

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \exists \varphi$$

CTL* path formulas

$$\varphi ::= \Phi \mid \varphi_1 \wedge \varphi_2 \mid \neg \varphi \mid \bigcirc \varphi \mid \varphi_1 \mathbf{U} \varphi_2$$

CTL: sublogic of **CTL***

- with path quantifiers $\exists$ and $\forall$
- restricted syntax of **path formulas**:

CTL* state formulas

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \exists \varphi$$

CTL* path formulas

$$\varphi ::= \Phi \mid \varphi_1 \wedge \varphi_2 \mid \neg \varphi \mid \bigcirc \varphi \mid \varphi_1 \mathbf{U} \varphi_2$$

CTL: sublogic of **CTL***

- with path quantifiers $\exists$ and $\forall$
- restricted syntax of **path formulas**:
 - * *no* boolean combinations of path formulas
 - * arguments of temporal operators $\bigcirc$ and $\mathbf{U}$ are **state formulas**

Let s_1, s_2 be states of a TS $\mathcal{T}$ without terminal states

Let s_1, s_2 be states of a TS $\mathcal{T}$ without terminal states


s_1, s_2 are **CTL** equivalent if for all **CTL** formulas ϕ :

$$s_1 \models \phi \quad \text{iff} \quad s_2 \models \phi$$

Let s_1, s_2 be states of a TS $\mathcal{T}$ without terminal states

s_1, s_2 are **CTL** equivalent if for all **CTL** formulas ϕ :


$$s_1 \models \phi \quad \text{iff} \quad s_2 \models \phi$$



Let s_1, s_2 be states of a TS $\mathcal{T}$ without terminal states

s_1, s_2 are **CTL** equivalent if for all **CTL** formulas ϕ :

$$s_1 \models \phi \quad \text{iff} \quad s_2 \models \phi$$



s_1, s_2 are
not **CTL** equivalent

$$s_1 \models \exists O(\exists O(\exists O \textcolor{red}{a} \vee \exists O \textcolor{green}{b}))$$

$$s_2 \not\models \exists O(\exists O(\exists O \textcolor{red}{a} \vee \exists O \textcolor{green}{b}))$$

Let s_1, s_2 be states of a TS $\mathcal{T}$ without terminal states

s_1, s_2 are **CTL** equivalent if for all **CTL** formulas ϕ :

$$s_1 \models \phi \quad \text{iff} \quad s_2 \models \phi$$

analogous definition for **CTL*** and **LTL**

Let s_1, s_2 be states of a TS $\mathcal{T}$ without terminal states

s_1, s_2 are **CTL** equivalent if for all **CTL** formulas ϕ :

$$s_1 \models \phi \quad \text{iff} \quad s_2 \models \phi$$

s_1, s_2 are **CTL*** equivalent if for all **CTL*** formulas ϕ :

$$s_1 \models \phi \quad \text{iff} \quad s_2 \models \phi$$

s_1, s_2 are **LTL** equivalent if for all **LTL** formulas φ :

$$s_1 \models \varphi \quad \text{iff} \quad s_2 \models \varphi$$

bisimulation equivalence
= CTL equivalence
= CTL* equivalence

bisimulation equivalence
= CTL equivalence
= CTL* equivalence

← for finite TS

bisimulation equivalence
= CTL equivalence
= CTL* equivalence


← for finite TS


Let $\mathcal{T}$ be a finite TS without terminal states,
and s_1, s_2 states in $\mathcal{T}$. Then:


$$s_1 \sim_{\mathcal{T}} s_2$$


iff s_1 and s_2 are CTL equivalent

iff s_1 and s_2 are CTL* equivalent









For arbitrary (possibly infinite) transition systems without terminal states:

For arbitrary (possibly infinite) transition systems without terminal states:

If s_1, s_2 are states with $s_1 \sim_{\mathcal{T}} s_2$ then for all CTL* formulas Φ :

$$s_1 \models \Phi \quad \text{iff} \quad s_2 \models \Phi$$

show by structural induction on CTL* formulas:

- (a) if s_1, s_2 are states with $s_1 \sim_{\mathcal{T}} s_2$ then
for all CTL* state formulas Φ :

$$s_1 \models \Phi \text{ iff } s_2 \models \Phi$$

- (b) if π_1, π_2 are paths with $\pi_1 \sim_{\mathcal{T}} \pi_2$ then
for all CTL* path formulas φ :

$$\pi_1 \models \varphi \text{ iff } \pi_2 \models \varphi$$

show by structural induction on CTL* formulas:

- (a) if s_1, s_2 are states with $s_1 \sim_T s_2$ then
for all CTL* state formulas Φ :

$$s_1 \models \Phi \text{ iff } s_2 \models \Phi$$

- (b) if π_1, π_2 are paths with $\pi_1 \sim_T \pi_2$ then
for all CTL* path formulas φ :


$$\pi_1 \models \varphi \text{ iff } \pi_2 \models \varphi$$

$\pi_1 \sim_T \pi_2 \stackrel{\text{def}}{\iff} \pi_1 \text{ and } \pi_2 \text{ are statewise bisimulation equivalent}$

Bisimulation equivalence $\Rightarrow$ CTL* equivalence

CTLEQ5.2-3

statewise bisimulation equivalent paths:



Bisimulation equivalence $\Rightarrow$ CTL* equivalence

CTLEQ5.2-5

For all CTL* state formulas Φ and path formulas φ :

(a) if $s_1 \sim_{\mathcal{T}} s_2$ then: $s_1 \models \Phi$ iff $s_2 \models \Phi$

(b) if $\pi_1 \sim_{\mathcal{T}} \pi_2$ then: $\pi_1 \models \varphi$ iff $\pi_2 \models \varphi$

Bisimulation equivalence $\Rightarrow$ CTL* equivalence

CTLEQ5.2-5

For all CTL* state formulas Φ and path formulas φ :

(a) if $s_1 \sim_{\mathcal{T}} s_2$ then: $s_1 \models \Phi$ iff $s_2 \models \Phi$

(b) if $\pi_1 \sim_{\mathcal{T}} \pi_2$ then: $\pi_1 \models \varphi$ iff $\pi_2 \models \varphi$

Proof by structural induction

Bisimulation equivalence $\Rightarrow$ CTL* equivalence

CTLEQ5.2-5

For all CTL* state formulas Φ and path formulas φ :

(a) if $s_1 \sim_{\mathcal{T}} s_2$ then: $s_1 \models \Phi$ iff $s_2 \models \Phi$

(b) if $\pi_1 \sim_{\mathcal{T}} \pi_2$ then: $\pi_1 \models \varphi$ iff $\pi_2 \models \varphi$

Proof by structural induction

base of induction:

(a) $\Phi = \text{true}$ or $\Phi = a \in AP$

(b) $\varphi = \Phi$ for some state formula Φ
s.t. statement (a) holds for Φ

Bisimulation equivalence $\Rightarrow$ CTL* equivalence

CTLEQ5.2-5

For all CTL* state formulas Φ and path formulas φ :

(a) if $s_1 \sim_T s_2$ then: $s_1 \models \Phi$ iff $s_2 \models \Phi$

(b) if $\pi_1 \sim_T \pi_2$ then: $\pi_1 \models \varphi$ iff $\pi_2 \models \varphi$

Proof by structural induction

step of induction:

(a) consider $\Phi = \Phi_1 \wedge \Phi_2, \neg\Psi$ or $\exists\varphi$ s.t.

(a) holds for Φ_1, Φ_2, Ψ


(b) holds for φ

(b) consider $\varphi = \varphi_1 \wedge \varphi_2, \neg\varphi', \bigcirc\varphi', \varphi_1 \mathbf{U} \varphi_2$ s.t.

(b) holds for $\varphi_1, \varphi_2, \varphi'$


Path lifting for $\sim_{\mathcal{T}}$

CTLEQ5.2-4



can be
completed to





If $s_1 \sim_{\mathcal{T}} s_2$ then for all $\pi_1 \in \text{Paths}(s_1)$
there exists $\pi_2 \in \text{Paths}(s_2)$ with $\pi_1 \sim_{\mathcal{T}} \pi_2$

$s_1 \sim_{\mathcal{T}} s_2$

$\downarrow$

s_{11}

$\downarrow$

s_{21}

$\downarrow$

s_{31}

$\downarrow$

$\vdots$

path π_1

can be
completed to

$s_1 \sim_{\mathcal{T}} s_2$

$\downarrow$

s_{11}

$\downarrow$

s_{21}

$\downarrow$

s_{31}

$\downarrow$

$\vdots$

$\sim_{\mathcal{T}}$

$\sim_{\mathcal{T}}$

$\sim_{\mathcal{T}}$

$\sim_{\mathcal{T}}$

s_2

$\downarrow$

s_{12}

$\downarrow$

s_{22}


$\downarrow$

s_{32}

$\downarrow$

$\vdots$

If $s_1 \sim_{\mathcal{T}} s_2$ then for all $\pi_1 \in \text{Paths}(s_1)$
there exists $\pi_2 \in \text{Paths}(s_2)$ with $\pi_1 \sim_{\mathcal{T}} \pi_2$



If $s_1 \sim_{\mathcal{T}} s_2$ then for all $\pi_1 \in \text{Paths}(s_1)$
there exists $\pi_2 \in \text{Paths}(s_2)$ with $\pi_1 \sim_{\mathcal{T}} \pi_2$

Correct or wrong?

CTLEQ5.2-6

If s_1, s_2 are not **CTL** equivalent then there exists a **CTL** formula Φ with $s_1 \models \Phi$ and $s_2 \not\models \Phi$

Correct or wrong?

CTLEQ5.2-6

If s_1, s_2 are not CTL equivalent then there exists a CTL formula ϕ with $s_1 \models \phi$ and $s_2 \not\models \phi$

correct.

If s_1, s_2 are not **CTL** equivalent then there exists a **CTL** formula Φ with $s_1 \models \Phi$ and $s_2 \not\models \Phi$

correct.

If s_1, s_2 not **CTL** equivalent then
there exists a **CTL** formula Φ with

$$s_1 \models \Phi \wedge s_2 \not\models \Phi$$

or $s_1 \not\models \Phi \wedge s_2 \models \Phi$

If s_1, s_2 are not **CTL** equivalent then there exists a **CTL** formula Φ with $s_1 \models \Phi$ and $s_2 \not\models \Phi$

correct.

If s_1, s_2 not **CTL** equivalent then
there exists a **CTL** formula Φ with

$$s_1 \models \Phi \wedge s_2 \not\models \Phi$$

$$\text{or } s_1 \not\models \Phi \wedge s_2 \models \Phi \implies s_1 \models \neg\Phi \wedge s_2 \not\models \neg\Phi$$

If s_1, s_2 are not **CTL** equivalent then there exists a **CTL** formula ϕ with $s_1 \models \phi$ and $s_2 \not\models \phi$

correct.

If s_1, s_2 are not **LTL** equivalent then there exists a **LTL** formula φ with $s_1 \models \varphi$ and $s_2 \not\models \varphi$

If s_1, s_2 are not **CTL** equivalent then there exists a **CTL** formula ϕ with $s_1 \models \phi$ and $s_2 \not\models \phi$

correct.

If s_1, s_2 are not **LTL** equivalent then there exists a **LTL** formula φ with $s_1 \models \varphi$ and $s_2 \not\models \varphi$


wrong.

If s_1, s_2 are not **CTL** equivalent then there exists a **CTL** formula Φ with $s_1 \models \Phi$ and $s_2 \not\models \Phi$

correct.

If s_1, s_2 are not **LTL** equivalent then there exists a **LTL** formula φ with $s_1 \models \varphi$ and $s_2 \not\models \varphi$

wrong.




If s_1, s_2 are not **CTL** equivalent then there exists a **CTL** formula Φ with $s_1 \models \Phi$ and $s_2 \not\models \Phi$

correct.

If s_1, s_2 are not **LTL** equivalent then there exists a **LTL** formula φ with $s_1 \models \varphi$ and $s_2 \not\models \varphi$

wrong.

$$\text{Traces}(s_2) \subset \text{Traces}(s_1)$$



Correct or wrong?

CTLEQ5.2-6

If s_1, s_2 are not **CTL** equivalent then there exists a **CTL** formula Φ with $s_1 \models \Phi$ and $s_2 \not\models \Phi$


correct.

If s_1, s_2 are not **LTL** equivalent then there exists a **LTL** formula φ with $s_1 \models \varphi$ and $s_2 \not\models \varphi$

wrong.

$Traces(s_2) \subset Traces(s_1)$

hence: $s_1 \models \varphi$ implies $s_2 \models \varphi$



CTL equivalence $\Rightarrow$ bisimulation equivalence

CTLEQ5.2-7A

CTL equivalence $\implies$ bisimulation equivalence

CTLEQ5.2-7A

If $\mathcal{T}$ is a finite TS then, for all states s_1, s_2 in $\mathcal{T}$:
if s_1, s_2 are **CTL** equivalent then $s_1 \sim_{\mathcal{T}} s_2$

CTL equivalence $\implies$ bisimulation equivalence

CTLEQ5.2-7A

If $\mathcal{T}$ is a **finite** TS then, for all states s_1, s_2 in $\mathcal{T}$:
if s_1, s_2 are **CTL** equivalent then $s_1 \sim_{\mathcal{T}} s_2$

CTL equivalence $\implies$ bisimulation equivalence

CTLEQ5.2-7A

If $\mathcal{T}$ is a **finite** TS then, for all states s_1, s_2 in $\mathcal{T}$:
if s_1, s_2 are **CTL** equivalent then $s_1 \sim_{\mathcal{T}} s_2$

Proof: show that

$\mathcal{R} \stackrel{\text{def}}{=} \{ (s_1, s_2) : s_1, s_2 \text{ satisfy the same } \mathbf{CTL} \text{ formulas} \}$

is a bisimulation

If $\mathcal{T}$ is a **finite** TS then, for all states s_1, s_2 in $\mathcal{T}$:
if s_1, s_2 are **CTL** equivalent then $s_1 \sim_{\mathcal{T}} s_2$

Proof: show that


$\mathcal{R} \stackrel{\text{def}}{=} \{ (s_1, s_2) : s_1, s_2 \text{ satisfy the same CTL formulas} \}$

is a bisimulation, i.e., for all $(s_1, s_2) \in \mathcal{R}$:

- (1) $L(s_1) = L(s_2)$
- (2) if $s_1 \rightarrow t_1$ then there exists a transition $s_2 \rightarrow t_2$
s.t. $(t_1, t_2) \in \mathcal{R}$

Example: CTL master formulas

CTLEQ5.2-7




$\bullet \hat{=} \{a\}$

$\bullet \hat{=} \{b\}$

$\bullet \hat{=} \emptyset$

Example: CTL master formulas

CTLEQ5.2-7



bisimulation equivalence $\sim_{\mathcal{T}}$
 $= \{ (v_1, v_2), (w_1, w_2), \dots \}$


$\bullet \equiv \{a\}$

$\bullet \equiv \{b\}$

$\bullet \equiv \emptyset$

Example: CTL master formulas

CTLEQ5.2-7



$\bullet \text{ (blue)} \hat{=} \{a\}$

$\bullet \text{ (red)} \hat{=} \{b\}$



$\bullet \text{ (green)} \hat{=} \emptyset$

bisimulation equivalence $\sim_{\mathcal{T}}$
 $= \{(v_1, v_2), (w_1, w_2), \dots\}$

but $u_1 \not\sim_{\mathcal{T}} u_2$

Example: CTL master formulas

CTLEQ5.2-7



bisimulation equivalence $\sim_{\mathcal{T}}$
 $= \{(v_1, v_2), (w_1, w_2), \dots\}$


but $u_1 \not\sim_{\mathcal{T}} u_2$

as $u_1 \rightarrow \{w_1, w_2\}$

$u_2 \not\rightarrow \{w_1, w_2\}$

Example: CTL master formulas

CTLEQ5.2-7



$\bullet \equiv \{a\}$

$\bullet \equiv \{b\}$

$\bullet \equiv \emptyset$

bisimulation equivalence $\sim_{\mathcal{T}}$
 $= \{(v_1, v_2), (w_1, w_2), \dots\}$

CTL master formulas:

$w_1, w_2 \models ?$



$v_1, v_2 \models ?$

$u_1 \models ?$

$u_2 \models ?$

Example: CTL master formulas

CTLEQ5.2-7



bisimulation equivalence $\sim_{\mathcal{T}}$
 $= \{(v_1, v_2), (w_1, w_2), \dots\}$

CTL master formulas:

$w_1, w_2 \models b$



$v_1, v_2 \models ?$

$u_1 \models ?$

$u_2 \models ?$

Example: CTL master formulas

CTLEQ5.2-7



bisimulation equivalence $\sim_{\mathcal{T}}$
 $= \{(v_1, v_2), (w_1, w_2), \dots\}$

CTL master formulas:

$$w_1, w_2 \models b$$


$$v_1, v_2 \models \neg a \wedge \neg b$$

$$u_1 \models ?$$

$$u_2 \models ?$$

Example: CTL master formulas

CTLEQ5.2-7



$$\text{blue node} \hat{=} \{a\}$$

$$\text{red node} \hat{=} \{b\}$$

$$\text{green node} \hat{=} \emptyset$$

bisimulation equivalence $\sim_{\mathcal{T}}$
 $= \{ (v_1, v_2), (w_1, w_2), \dots \}$

CTL master formulas:

$$w_1, w_2 \models b$$


$$v_1, v_2 \models \neg a \wedge \neg b$$

$$u_1 \models (\exists \bigcirc b) \wedge a$$

$$u_2 \models ?$$

Example: CTL master formulas

CTLEQ5.2-7



$$\text{blue node} \hat{=} \{a\}$$

$$\text{red node} \hat{=} \{b\}$$

$$\text{green node} \hat{=} \emptyset$$

bisimulation equivalence $\sim_{\mathcal{T}}$
 $= \{ (v_1, v_2), (w_1, w_2), \dots \}$

CTL master formulas:

$$w_1, w_2 \models b$$


$$v_1, v_2 \models \neg a \wedge \neg b$$

$$u_1 \models (\exists \bigcirc b) \wedge a$$

$$u_2 \models (\neg \exists \bigcirc b) \wedge a$$

...master formulas for $\sim_{\mathcal{T}}$ -classes?


CTLEQ5.2-8



$$AP = \{ \text{blue}, \text{red} \}$$

...master formulas for $\sim_T$ -classes?

CTLEQ5.2-8




$$AP = \{ \text{blue}, \text{red} \}$$

$$s_1 \sim_T s_2 \not\sim_T u$$

...master formulas for $\sim_T$ -classes?

CTLEQ5.2-8



$$AP = \{ \text{blue}, \text{red} \}$$

$$s_1 \sim_T s_2 \not\sim_T u$$

$$\phi_w = ?$$


$$\phi_C = ?$$

$$\phi_u = ?$$

$$\text{where } C = \{s_1, s_2\}$$

...master formulas for $\sim_T$ -classes?

CTLEQ5.2-8



$$AP = \{ \text{blue}, \text{red} \}$$

$$s_1 \sim_T s_2 \not\sim_T u$$

$$\Phi_w = \text{red}$$


$$\Phi_C = ?$$

$$\Phi_u = ?$$

$$\text{where } C = \{s_1, s_2\}$$

...master formulas for $\sim_T$ -classes?

CTLEQ5.2-8




$$AP = \{ \text{blue}, \text{red} \}$$

$$s_1 \sim_T s_2 \not\sim_T u$$

$$\Phi_w = \text{red}$$

$$\Phi_C = \text{blue} \wedge \forall \bigcirc \text{blue} \quad \text{where } C = \{s_1, s_2\}$$

$$\Phi_u = ?$$



$$AP = \{ \text{blue}, \text{red} \}$$

$$s_1 \sim_T s_2 \not\sim_T u$$

$$\Phi_w = \text{red}$$

$$\Phi_C = \text{blue} \wedge \forall \bigcirc \text{blue} \quad \text{where } C = \{s_1, s_2\}$$

$$\Phi_u = \exists \bigcirc \text{red}$$

CTL equivalence $\implies$ bisimulation equivalence

CTLEQ5.2-7B

If $\mathcal{T}$ is a finite TS then, for all states s_1, s_2 in $\mathcal{T}$:
if s_1, s_2 are **CTL** equivalent then $s_1 \sim_{\mathcal{T}} s_2$

CTL equivalence $\implies$ bisimulation equivalence

CTLEQ5.2-7B

If $\mathcal{T}$ is a **finite** TS then, for all states s_1, s_2 in $\mathcal{T}$:
if s_1, s_2 are **CTL** equivalent then $s_1 \sim_{\mathcal{T}} s_2$

- wrong for **infinite** TS

CTL equivalence $\implies$ bisimulation equivalence

CTLEQ5.2-7B

If $\mathcal{T}$ is a **finite** TS then, for all states s_1, s_2 in $\mathcal{T}$:
if s_1, s_2 are **CTL** equivalent then $s_1 \sim_{\mathcal{T}} s_2$

- wrong for **infinite** TS
- but also holds for **finitely branching** TS

CTL equivalence $\implies$ bisimulation equivalence

CTLEQ5.2-7B

If $\mathcal{T}$ is a **finite** TS then, for all states s_1, s_2 in $\mathcal{T}$:
if s_1, s_2 are **CTL** equivalent then $s_1 \sim_{\mathcal{T}} s_2$

- wrong for **infinite** TS
- but also holds for **finitely branching** TS



possibly infinite-state TS such that

- * the number of **initial states** is **finite**
- * for each state the number of **successors** is **finite**

CTL equivalence $\implies$ bisimulation equivalence


CTLEQ5.2-7C

Let $\mathcal{T} = (\mathcal{S}, Act, \rightarrow, \mathcal{S}_0, AP, L)$ be **finitely branching**.

CTL equivalence $\implies$ bisimulation equivalence

CTLEQ5.2-7c

Let $\mathcal{T} = (S, Act, \rightarrow, S_0, AP, L)$ be finitely branching.

- 
- * S_0 is finite
 - * $Post(s)$ is finite for all $s \in S$

CTL equivalence $\implies$ bisimulation equivalence

CTLEQ5.2-7C

Let $\mathcal{T} = (\mathcal{S}, Act, \rightarrow, \mathcal{S}_0, AP, L)$ be finitely branching.

- * $\mathcal{S}_0$ is finite
- * $Post(s)$ is finite for all $s \in \mathcal{S}$

Then, for all states s_1, s_2 in $\mathcal{T}$:

if s_1, s_2 are **CTL** equivalent then $s_1 \sim_{\mathcal{T}} s_2$

CTL equivalence $\implies$ bisimulation equivalence

CTLEQ5.2-7C

Let $\mathcal{T} = (\mathcal{S}, \text{Act}, \rightarrow, \mathcal{S}_0, \text{AP}, L)$ be finitely branching.

- * $\mathcal{S}_0$ is finite
- * $\text{Post}(s)$ is finite for all $s \in \mathcal{S}$

Then, for all states s_1, s_2 in $\mathcal{T}$:

if s_1, s_2 are **CTL** equivalent then $s_1 \sim_{\mathcal{T}} s_2$

Proof: as for finite TS.

CTL equivalence $\implies$ bisimulation equivalence

CTLEQ5.2-7c

Let $\mathcal{T} = (\mathcal{S}, Act, \rightarrow, \mathcal{S}_0, AP, L)$ be finitely branching.

- * $\mathcal{S}_0$ is finite
- * $Post(s)$ is finite for all $s \in \mathcal{S}$

Then, for all states s_1, s_2 in $\mathcal{T}$:

if s_1, s_2 are **CTL** equivalent then $s_1 \sim_{\mathcal{T}} s_2$

Proof: as for finite TS. Amounts showing that

$\mathcal{R} \stackrel{\text{def}}{=} \{ (s_1, s_2) : s_1, s_2 \text{ satisfy the same } \mathbf{CTL} \text{ formulas} \}$

is a bisimulation.

CTL equivalence $\implies$ bisimulation equivalence

CTLEQ5.2-7D

If $\mathcal{T}$ is a **finitely branching** TS then for all states s_1, s_2 :
if s_1, s_2 are **CTL** equivalent then $s_1 \sim_{\mathcal{T}} s_2$

Proof: show that

$\mathcal{R} \stackrel{\text{def}}{=} \{ (s_1, s_2) : s_1, s_2 \text{ satisfy the same CTL formulas} \}$

is a bisimulation, i.e., for $(s_1, s_2) \in \mathcal{R}$:

- (1) $L(s_1) = L(s_2)$
- (2) if $s_1 \rightarrow t_1$ then there exists a transition $s_2 \rightarrow t_2$
s.t. $(t_1, t_2) \in \mathcal{R}$

Let $\mathcal{T}$ be a **finite** TS without terminal states,
and s_1, s_2 states in $\mathcal{T}$. Then:

$$s_1 \sim_{\mathcal{T}} s_2$$

iff s_1 and s_2 are **CTL** equivalent


iff s_1 and s_2 are **CTL*** equivalent


Let $\mathcal{T}$ be a **finitely branching** TS without terminal states, and s_1, s_2 states in $\mathcal{T}$. Then:


$$s_1 \sim_{\mathcal{T}} s_2$$


iff s_1 and s_2 are **CTL** equivalent


iff s_1 and s_2 are **CTL*** equivalent











so far: we considered

- **CTL/CTL*** equivalence
- bisimulation equivalence $\sim_{\mathcal{T}}$

for the **states** of a single transition system $\mathcal{T}$

If $\mathcal{T}_1$, $\mathcal{T}_2$ are finitely branching TS over AP without terminal states then:

$$\mathcal{T}_1 \sim \mathcal{T}_2$$

iff $\mathcal{T}_1$ and $\mathcal{T}_2$ satisfy the same **CTL** formulas

iff $\mathcal{T}_1$ and $\mathcal{T}_2$ satisfy the same **CTL*** formulas

Does the following statements hold for **finite TS** without terminal states ?

CTL equivalence is finer than **LTL** equivalence

CTL equivalence is finer than **LTL** equivalence

correct.

CTL equivalence is finer than **LTL** equivalence

correct.



CTL equivalence = **CTL*** equivalence

LTL is sublogic of **CTL***

Correct or wrong?

CTLEQ5.2-9

CTL equivalence is finer than **LTL** equivalence

correct.

LTL equivalence is finer than **CTL** equivalence

CTL equivalence is finer than **LTL** equivalence

correct.

LTL equivalence is finer than **CTL** equivalence


wrong.

CTL equivalence is finer than **LTL** equivalence

correct.

LTL equivalence is finer than **CTL** equivalence

wrong.




CTL equivalence is finer than **LTL** equivalence

correct.

LTL equivalence is finer than **CTL** equivalence

wrong.




s_1 , s_2 are trace equivalent

CTL equivalence is finer than **LTL** equivalence

correct.

LTL equivalence is finer than **CTL** equivalence

wrong.




s_1 , s_2 are trace equivalent
and **LTL** equivalent

CTL equivalence is finer than **LTL** equivalence

correct.


LTL equivalence is finer than **CTL** equivalence


wrong.





s_1, s_2 are trace equivalent
and **LTL** equivalent

$$\begin{aligned} s_1 &\models \exists O(\exists O a \vee \exists O b) \\ s_2 &\not\models \exists O(\exists O a \vee \exists O b) \end{aligned}$$









Correct or wrong?

CTLEQ5.2-11

Let $\mathcal{T}$ be a finite TS without terminal states and s_1, s_2 states of $\mathcal{T}$.

If s_1, s_2 satisfy the same $\text{CTL}_{\setminus U}$ formulas then
 $s_1 \sim_{\mathcal{T}} s_2$.

Let $\mathcal{T}$ be a finite TS without terminal states and s_1, s_2 states of $\mathcal{T}$.

If s_1, s_2 satisfy the same $\text{CTL}_{\setminus \mathbf{U}}$ formulas then
 $s_1 \sim_{\mathcal{T}} s_2$.

where $\text{CTL}_{\setminus \mathbf{U}} \hat{=} \text{CTL}$ without until operator $\mathbf{U}$

Correct or wrong?

CTLEQ5.2-11

Let $\mathcal{T}$ be a finite TS without terminal states and s_1, s_2 states of $\mathcal{T}$.

If s_1, s_2 satisfy the same $\text{CTL} \setminus \mathbf{U}$ formulas then
 $s_1 \sim_{\mathcal{T}} s_2$.

where $\text{CTL} \setminus \mathbf{U} \hat{=} \text{CTL}$ without until operator $\mathbf{U}$

correct.

Let $\mathcal{T}$ be a finite TS without terminal states and s_1, s_2 states of $\mathcal{T}$.

If s_1, s_2 satisfy the same $\text{CTL}_{\setminus \mathbf{U}}$ formulas then
 $s_1 \sim_{\mathcal{T}} s_2$.

where $\text{CTL}_{\setminus \mathbf{U}} \hat{=} \text{CTL}$ without until operator $\mathbf{U}$

correct. see the proof

“ $\text{CTL equivalence} \implies \text{bisimulation equivalence}$ ”

$\text{CTL}_{\setminus U}$ -equivalence $\Rightarrow$ bisimulation equivalence CTLEQ5.2-11

Let $\mathcal{T}$ be a finite TS without terminal states and s_1, s_2 states of $\mathcal{T}$.

If s_1, s_2 satisfy the same $\text{CTL}_{\setminus U}$ formulas then
 $s_1 \sim_{\mathcal{T}} s_2$.

Proof. Show that $\text{CTL}_{\setminus U}$ equivalence is a bisimulation

$\text{CTL}_{\setminus U}$ -equivalence $\Rightarrow$ bisimulation equivalence CTLEQ5.2-11

Let $\mathcal{T}$ be a finite TS without terminal states and s_1, s_2 states of $\mathcal{T}$.

If s_1, s_2 satisfy the same $\text{CTL}_{\setminus U}$ formulas then
 $s_1 \sim_{\mathcal{T}} s_2$.

Proof. Show that $\text{CTL}_{\setminus U}$ equivalence is a bisimulation

- labeling condition only uses atomic propositions

CTL_{\U}-equivalence $\Rightarrow$ bisimulation equivalence CTLEQ5.2-11

Let $\mathcal{T}$ be a finite TS without terminal states and s_1, s_2 states of $\mathcal{T}$.

If s_1, s_2 satisfy the same CTL_{\U} formulas then
 $s_1 \sim_{\mathcal{T}} s_2$.

Proof. Show that CTL_{\U} equivalence is a bisimulation

- labeling condition only uses atomic propositions
- simulation condition can be established by CTL_{\U} master formulas of the form:

$\text{CTL}_{\setminus U}$ -equivalence $\Rightarrow$ bisimulation equivalence CTLEQ5.2-11

Let $\mathcal{T}$ be a finite TS without terminal states and s_1, s_2 states of $\mathcal{T}$.

If s_1, s_2 satisfy the same $\text{CTL}_{\setminus U}$ formulas then
 $s_1 \sim_{\mathcal{T}} s_2$.

Proof. Show that $\text{CTL}_{\setminus U}$ equivalence is a bisimulation

- labeling condition only uses atomic propositions
- simulation condition can be established by $\text{CTL}_{\setminus U}$ master formulas of the form:

$$\exists \bigcirc \Phi_C \quad \text{where} \quad \Phi_C = \bigwedge_D \Phi_{C,D}$$

$\text{CTL}_{\setminus U}$ -equivalence $\Rightarrow$ bisimulation equivalence CTLEQ5.2-11

Let $\mathcal{T}$ be a finite TS without terminal states and s_1, s_2 states of $\mathcal{T}$.

If s_1, s_2 satisfy the same $\text{CTL}_{\setminus U}$ formulas then
 $s_1 \sim_{\mathcal{T}} s_2$.

Proof. Show that $\text{CTL}_{\setminus U}$ equivalence is a bisimulation

- labeling condition only uses atomic propositions
- simulation condition can be established by $\text{CTL}_{\setminus U}$ master formulas of the form:

$$\exists \bigcirc \Phi_C \quad \text{where} \quad \Phi_C = \bigwedge_D \Phi_{C,D}$$

and $\text{Sat}(\Phi_{C,D}) \subseteq C \setminus D$

Let $\mathcal{T}$ be a finite TS without terminal states.

$\mathcal{T}$ and its bisimulation quotient $\mathcal{T}/\sim$ satisfy the same **CTL*** formulas.

Let $\mathcal{T}$ be a finite TS without terminal states.

$\mathcal{T}$ and its bisimulation quotient $\mathcal{T}/\sim$ satisfy the same **CTL*** formulas.

correct.

Let $\mathcal{T}$ be a finite TS without terminal states.

$\mathcal{T}$ and its bisimulation quotient $\mathcal{T}/\sim$ satisfy the same CTL* formulas.

correct. Recall that $\mathcal{T} \sim \mathcal{T}/\sim$

Let $\mathcal{T}$ be a finite TS without terminal states.

$\mathcal{T}$ and its bisimulation quotient $\mathcal{T}/\sim$ satisfy the same CTL* formulas.

correct. Recall that $\mathcal{T} \sim \mathcal{T}/\sim$ as

$$\mathcal{R} = \{(s, [s]) : s \in S\}$$

is a bisimulation for $(\mathcal{T}, \mathcal{T}/\sim)$

here: $[s] = \sim_{\mathcal{T}}$ -equivalence class of state s