

Compiler Construction

Lecture 9: Syntax Analysis V ($LR(k)$ Grammars)

Thomas Noll

Lehrstuhl für Informatik 2
(Software Modeling and Verification)

RWTH Aachen University

noll@cs.rwth-aachen.de

<http://www-i2.informatik.rwth-aachen.de/i2/cc12/>

Summer Semester 2012

- 1 Repetition: Nondeterministic Bottom-Up Parsing
- 2 Nondeterministic Bottom-Up Parsing (cont.)
- 3 $LR(k)$ Grammars
- 4 $LR(0)$ Grammars
- 5 Examples of $LR(0)$ Conflicts

Approach:

- 1 Given $G \in CFG_{\Sigma}$, construct a **nondeterministic bottom-up parsing automaton** (NBA) which accepts $L(G)$ and which additionally computes corresponding (reversed) rightmost analyses
 - input alphabet: Σ
 - pushdown alphabet: X
 - output alphabet: $[p]$ (where $p := |P|$)
 - state set: omitted
 - transitions:
 - shift**: shifting input symbols onto the pushdown
 - reduce**: replacing the right-hand side of a production by its left-hand side (= inverse expansion steps)
- 2 Remove nondeterminism by allowing **lookahead** on the input:
 $G \in LR(k)$ iff $L(G)$ recognizable by deterministic bottom-up parsing automaton with lookahead of k symbols

Nondeterministic Bottom-Up Automaton

Definition (Nondeterministic bottom-up parsing automaton)

Let $G = \langle N, \Sigma, P, S \rangle \in CFG_{\Sigma}$. The **nondeterministic bottom-up parsing automaton** of G , $NBA(G)$, is defined by the following components.

- **Input alphabet:** Σ
- **Pushdown alphabet:** X
- **Output alphabet:** $[p]$
- **Configurations:** $\Sigma^* \times X^* \times [p]^*$ (top of pushdown to the right)
- **Transitions** for $w \in \Sigma^*$, $\alpha \in X^*$, and $z \in [p]^*$:
 - shifting steps: $(aw, \alpha, z) \vdash (w, \alpha a, z)$ if $a \in \Sigma$
 - reduction steps: $(w, \alpha\beta, z) \vdash (w, \alpha A, zi)$ if $\pi_i = A \rightarrow \beta$
- **Initial configuration** for $w \in \Sigma^*$: $(w, \varepsilon, \varepsilon)$
- **Final configurations:** $\{\varepsilon\} \times \{S\} \times [p]^*$

- 1 Repetition: Nondeterministic Bottom-Up Parsing
- 2 Nondeterministic Bottom-Up Parsing (cont.)
- 3 $LR(k)$ Grammars
- 4 $LR(0)$ Grammars
- 5 Examples of $LR(0)$ Conflicts

Theorem 9.1 (Correctness of $\text{NBA}(G)$)

Let $G = \langle N, \Sigma, P, S \rangle \in \text{CFG}_\Sigma$ and $\text{NBA}(G)$ as before. Then, for every $w \in \Sigma^*$ and $z \in [p]^*$,

$(w, \varepsilon, \varepsilon) \vdash^* (\varepsilon, S, z)$ iff $\overleftarrow{z}$ is a rightmost analysis of w

Theorem 9.1 (Correctness of $\text{NBA}(G)$)

Let $G = \langle N, \Sigma, P, S \rangle \in \text{CFG}_\Sigma$ and $\text{NBA}(G)$ as before. Then, for every $w \in \Sigma^*$ and $z \in [p]^*$,

$(w, \varepsilon, \varepsilon) \vdash^* (\varepsilon, S, z)$ iff $\overleftarrow{z}$ is a rightmost analysis of w

Proof.

similar to the top-down case (Theorem 6.1)



Nondeterminism in $\text{NBA}(G)$

Observation: $\text{NBA}(G)$ is generally **nondeterministic**

- **Shift or reduce?** Example:

$$(bw, \alpha a, z) \vdash \begin{cases} (w, \alpha ab, z) \\ (bw, \alpha A, zi) \end{cases} \text{ if } \pi_i = A \rightarrow a$$

Nondeterminism in NBA(G)

Observation: NBA(G) is generally **nondeterministic**

- **Shift or reduce?** Example:

$$(bw, \alpha a, z) \vdash \begin{cases} (w, \alpha ab, z) \\ (bw, \alpha A, zi) \end{cases} \text{ if } \pi_i = A \rightarrow a$$

- If reduce: **which “handle” β ?** Example:

$$(w, \alpha ab, z) \vdash \begin{cases} (w, \alpha A, zi) \\ (w, \alpha aB, zj) \end{cases} \text{ if } \pi_i = A \rightarrow ab \text{ and } \pi_j = B \rightarrow b$$

Nondeterminism in $NBA(G)$

Observation: $NBA(G)$ is generally **nondeterministic**

- **Shift or reduce?** Example:

$$(bw, \alpha a, z) \vdash \begin{cases} (w, \alpha ab, z) \\ (bw, \alpha A, zi) \end{cases} \text{ if } \pi_i = A \rightarrow a$$

- If reduce: **which “handle” β ?** Example:

$$(w, \alpha ab, z) \vdash \begin{cases} (w, \alpha A, zi) \\ (w, \alpha aB, zj) \end{cases} \text{ if } \pi_i = A \rightarrow ab \text{ and } \pi_j = B \rightarrow b$$

- If reduce β : **which left-hand side A ?** Example:

$$(w, \alpha a, z) \vdash \begin{cases} (w, \alpha A, zi) \\ (w, \alpha B, zj) \end{cases} \text{ if } \pi_i = A \rightarrow a \text{ and } \pi_j = B \rightarrow a$$

Nondeterminism in NBA(G)

Observation: NBA(G) is generally **nondeterministic**

- **Shift or reduce?** Example:

$$(bw, \alpha a, z) \vdash \begin{cases} (w, \alpha ab, z) \\ (bw, \alpha A, zi) \end{cases} \text{ if } \pi_i = A \rightarrow a$$

- If reduce: **which “handle” β ?** Example:

$$(w, \alpha ab, z) \vdash \begin{cases} (w, \alpha A, zi) \\ (w, \alpha aB, zj) \end{cases} \text{ if } \pi_i = A \rightarrow ab \text{ and } \pi_j = B \rightarrow b$$

- If reduce β : **which left-hand side A ?** Example:

$$(w, \alpha a, z) \vdash \begin{cases} (w, \alpha A, zi) \\ (w, \alpha B, zj) \end{cases} \text{ if } \pi_i = A \rightarrow a \text{ and } \pi_j = B \rightarrow a$$

- **When to terminate parsing?** Example:

$$\underbrace{(\varepsilon, S, z)}_{\text{final}} \vdash (\varepsilon, A, zi) \text{ if } \pi_i = A \rightarrow S$$

General assumption to avoid nondeterminism of last type:
every grammar is start separated

Definition 9.2 (Start separation)

A grammar $G = \langle N, \Sigma, P, S \rangle \in CFG_{\Sigma}$ is called **start separated** if S only occurs in productions of the form $S \rightarrow A$ where $A \neq S$.

General assumption to avoid nondeterminism of last type:
every grammar is start separated

Definition 9.2 (Start separation)

A grammar $G = \langle N, \Sigma, P, S \rangle \in CFG_{\Sigma}$ is called **start separated** if S only occurs in productions of the form $S \rightarrow A$ where $A \neq S$.

Remarks:

- Start separation always possible by adding $S' \rightarrow S$ with **new start symbol** S'
- From now on consider only **reduced** grammars of this form
($\pi_0 := S' \rightarrow S$)

Resolving Termination Nondeterminism II

Start separation removes last form of nondeterminism (“When to terminate parsing?”):

Lemma 9.3

*If $G \in CFG_{\Sigma}$ is start separated, then no successor of a final configuration (ϵ, S', z) in $NBA(G)$ is again a final configuration.
(Thus parsing should be stopped in the first final configuration.)*

Resolving Termination Nondeterminism II

Start separation removes last form of nondeterminism (“When to terminate parsing?”):

Lemma 9.3

*If $G \in CFG_{\Sigma}$ is start separated, then no successor of a final configuration (ε, S', z) in $NBA(G)$ is again a final configuration.
(Thus parsing should be stopped in the first final configuration.)*

Proof.

- To (ε, S', z) , only reductions by ε -productions can be applied:

$$(\varepsilon, S', z) \vdash (\varepsilon, S'A, zi) \quad \text{if } \pi_i = A \rightarrow \varepsilon$$

Resolving Termination Nondeterminism II

Start separation removes last form of nondeterminism (“When to terminate parsing?”):

Lemma 9.3

*If $G \in CFG_{\Sigma}$ is start separated, then no successor of a final configuration (ε, S', z) in $NBA(G)$ is again a final configuration.
(Thus parsing should be stopped in the first final configuration.)*

Proof.

- To (ε, S', z) , only reductions by ε -productions can be applied:
$$(\varepsilon, S', z) \vdash (\varepsilon, S' A, zi) \quad \text{if } \pi_i = A \rightarrow \varepsilon$$
- Thereafter, only reductions by productions of the form $A_0 \rightarrow A_1 \dots A_n$ ($n \geq 0$) can be applied

Resolving Termination Nondeterminism II

Start separation removes last form of nondeterminism (“When to terminate parsing?”):

Lemma 9.3

*If $G \in CFG_{\Sigma}$ is start separated, then no successor of a final configuration (ε, S', z) in $NBA(G)$ is again a final configuration.
(Thus parsing should be stopped in the first final configuration.)*

Proof.

- To (ε, S', z) , only reductions by ε -productions can be applied:

$$(\varepsilon, S', z) \vdash (\varepsilon, S' A, zi) \quad \text{if } \pi_i = A \rightarrow \varepsilon$$

- Thereafter, only reductions by productions of the form $A_0 \rightarrow A_1 \dots A_n$ ($n \geq 0$) can be applied
- Every resulting configuration is of the (non-final) form

$$(\varepsilon, S' B_1 \dots B_k, z) \quad \text{where } k \geq 1$$



- 1 Repetition: Nondeterministic Bottom-Up Parsing
- 2 Nondeterministic Bottom-Up Parsing (cont.)
- 3 $LR(k)$ Grammars
- 4 $LR(0)$ Grammars
- 5 Examples of $LR(0)$ Conflicts

Goal: resolve remaining nondeterminism of $NBA(G)$ by supporting lookahead of $k \in \mathbb{N}$ symbols on the input

$\Rightarrow LR(k)$: reading of input from left to right with k -lookahead, computing a rightmost analysis

Goal: resolve remaining nondeterminism of $NBA(G)$ by supporting **lookahead of $k \in \mathbb{N}$ symbols** on the input

$\Rightarrow LR(k)$: reading of input from **l**eft to **r**ight with k -lookahead, computing a **r**ightmost analysis

Definition 9.4 ($LR(k)$ grammar)

Let $G = \langle N, \Sigma, P, S \rangle \in CFG_{\Sigma}$ be start separated and $k \in \mathbb{N}$. Then G has the **$LR(k)$ property** (notation: $G \in LR(k)$) if for all rightmost derivations of the form

$$S \begin{cases} \Rightarrow_r^* \alpha A w \Rightarrow_r \alpha \beta w \\ \Rightarrow_r^* \gamma B x \Rightarrow_r \alpha \beta y \end{cases}$$

such that $\text{first}_k(w) = \text{first}_k(y)$, it follows that $\alpha = \gamma$, $A = B$, and $x = y$.

Remarks:

- If $G \in LR(k)$, then the reduction of $\alpha\beta w$ to αAw is already determined by $\text{first}_k(w)$.
- Therefore $\text{NBA}(G)$ in configuration $(w, \alpha\beta, z)$ can decide to reduce and how to reduce.

Remarks:

- If $G \in LR(k)$, then the reduction of $\alpha\beta w$ to $\alpha A w$ is already determined by $\text{first}_k(w)$.
- Therefore $\text{NBA}(G)$ in configuration $(w, \alpha\beta, z)$ can decide to reduce and how to reduce.
- **Computation of $\text{NBA}(G)$** for $S \Rightarrow_r^* \alpha A w \Rightarrow_r \alpha\beta w$:

$$(w'w, \varepsilon, \varepsilon) \vdash^* (w, \alpha\beta, z) \stackrel{\text{red } i}{\vdash} (w, \alpha A, zi) \vdash \dots$$

where $\pi_i = A \rightarrow \beta$

Remarks:

- If $G \in LR(k)$, then the reduction of $\alpha\beta w$ to $\alpha A w$ is already determined by $\text{first}_k(w)$.
- Therefore $\text{NBA}(G)$ in configuration $(w, \alpha\beta, z)$ can decide to reduce and how to reduce.
- **Computation of $\text{NBA}(G)$** for $S \Rightarrow_r^* \alpha A w \Rightarrow_r \alpha\beta w$:

$$(w'w, \varepsilon, \varepsilon) \vdash^* (w, \alpha\beta, z) \xrightarrow{\text{red } i} (w, \alpha A, zi) \vdash \dots$$

where $\pi_i = A \rightarrow \beta$

- **Computation of $\text{NBA}(G)$** for $S \Rightarrow_r^* \gamma B x \Rightarrow_r \alpha\beta y$:
 - with direct reduction ($y = x$, $\alpha\beta = \gamma\delta$, $\pi_j = B \rightarrow \delta$):

$$(y'y, \varepsilon, \varepsilon) \vdash^* (y, \alpha\beta, z') \xrightarrow{\text{red } j} (x, \gamma B, z'j) \vdash \dots$$

Remarks:

- If $G \in LR(k)$, then the reduction of $\alpha\beta w$ to $\alpha A w$ is already determined by $\text{first}_k(w)$.
- Therefore $\text{NBA}(G)$ in configuration $(w, \alpha\beta, z)$ can decide to reduce and how to reduce.
- **Computation of $\text{NBA}(G)$** for $S \Rightarrow_r^* \alpha A w \Rightarrow_r \alpha\beta w$:

$$(w'w, \varepsilon, \varepsilon) \vdash^* (w, \alpha\beta, z) \xrightarrow{\text{red } i} (w, \alpha A, zi) \vdash \dots$$

where $\pi_i = A \rightarrow \beta$

- **Computation of $\text{NBA}(G)$** for $S \Rightarrow_r^* \gamma B x \Rightarrow_r \alpha\beta y$:
 - with direct reduction ($y = x$, $\alpha\beta = \gamma\delta$, $\pi_j = B \rightarrow \delta$):

$$(y'y, \varepsilon, \varepsilon) \vdash^* (y, \alpha\beta, z') \xrightarrow{\text{red } j} (x, \gamma B, z'j) \vdash \dots$$

- with previous shifts ($y = x'x$, $\alpha\beta x' = \gamma\delta$, $\pi_j = B \rightarrow \delta$):

$$\begin{aligned} (y'y, \varepsilon, \varepsilon) &\vdash^* (y, \alpha\beta, z') = (x'x, \alpha\beta, z') \\ &\xrightarrow{\text{shift}^*} (x, \alpha\beta x', z') = (x, \gamma\delta, z') \\ &\xrightarrow{\text{red } j} (x, \gamma B, z'j) \vdash \dots \end{aligned}$$

- 1 Repetition: Nondeterministic Bottom-Up Parsing
- 2 Nondeterministic Bottom-Up Parsing (cont.)
- 3 $LR(k)$ Grammars
- 4 $LR(0)$ Grammars
- 5 Examples of $LR(0)$ Conflicts

The case $k = 0$ is relevant (in contrast to $LL(0)$): here the decision is just based on the contents of the pushdown, **without any lookahead**.

The case $k = 0$ is relevant (in contrast to $LL(0)$): here the decision is just based on the contents of the pushdown, **without any lookahead**.

Corollary 9.5 ($LR(0)$ grammar)

$G \in CFG_{\Sigma}$ has the $LR(0)$ property if for all rightmost derivations of the form

$$S \begin{cases} \Rightarrow_r^* \alpha Aw \Rightarrow_r \alpha \beta w \\ \Rightarrow_r^* \gamma Bx \Rightarrow_r \alpha \beta y \end{cases}$$

it follows that $\alpha = \gamma$, $A = B$, and $x = y$.

The case $k = 0$ is relevant (in contrast to $LL(0)$): here the decision is just based on the contents of the pushdown, **without any lookahead**.

Corollary 9.5 ($LR(0)$ grammar)

$G \in CFG_{\Sigma}$ has the **$LR(0)$ property** if for all rightmost derivations of the form

$$S \begin{cases} \Rightarrow_r^* \alpha A w \Rightarrow_r \alpha \beta w \\ \Rightarrow_r^* \gamma B x \Rightarrow_r \alpha \beta y \end{cases}$$


it follows that $\alpha = \gamma$, $A = B$, and $x = y$.

Goal: derive a **finite information** from the pushdown which suffices to resolve the nondeterminism (similar to abstraction of right context in LL parsing by fo-sets)

Example 9.6

$G :$

$S' \rightarrow S$	(1)
$S \rightarrow B \mid C$	(2,3)
$B \rightarrow aB \mid b$	(4,5)
$C \rightarrow aC \mid c$	(6,7)




Example 9.6

$G :$

$S' \rightarrow S$	(1)
$S \rightarrow B \mid C$	(2,3)
$B \rightarrow aB \mid b$	(4,5)
$C \rightarrow aC \mid c$	(6,7)

NBA(G):
($ab, \varepsilon, \varepsilon$)




Example 9.6

$G :$

$S' \rightarrow S$	(1)
$S \rightarrow B \mid C$	(2,3)
$B \rightarrow aB \mid b$	(4,5)
$C \rightarrow aC \mid c$	(6,7)

NBA(G):
 $(ab, \varepsilon, \varepsilon)$
 $\vdash (b, a, \varepsilon)$




Example 9.6

$G :$

$S' \rightarrow S$	(1)
$S \rightarrow B \mid C$	(2,3)
$B \rightarrow aB \mid b$	(4,5)
$C \rightarrow aC \mid c$	(6,7)

NBA(G):

$(ab, \varepsilon, \varepsilon)$
 $\vdash (b, a, \varepsilon)$
 $\vdash (\varepsilon, ab, \varepsilon)$




Example 9.6

$G :$

$S' \rightarrow S$	(1)
$S \rightarrow B \mid C$	(2,3)
$B \rightarrow aB \mid b$	(4,5)
$C \rightarrow aC \mid c$	(6,7)

NBA(G):

$(ab, \varepsilon, \varepsilon)$
 $\vdash (b, a, \varepsilon)$
 $\vdash (\varepsilon, ab, \varepsilon)$
 $\vdash (\varepsilon, aB, 5)$




Example 9.6

$G :$

$S' \rightarrow S$	(1)
$S \rightarrow B \mid C$	(2,3)
$B \rightarrow aB \mid b$	(4,5)
$C \rightarrow aC \mid c$	(6,7)

NBA(G):

$(ab, \varepsilon, \varepsilon)$
 $\vdash (b, a, \varepsilon)$
 $\vdash (\varepsilon, ab, \varepsilon)$
 $\vdash (\varepsilon, aB, 5)$
 $\vdash (\varepsilon, B, 54)$




Example 9.6

$G :$ $S' \rightarrow S$ (1)
 $S \rightarrow B \mid C$ (2,3)
 $B \rightarrow aB \mid b$ (4,5)
 $C \rightarrow aC \mid c$ (6,7)

NBA(G):

$(ab, \varepsilon, \varepsilon)$
 $\vdash (b, a, \varepsilon)$
 $\vdash (\varepsilon, ab, \varepsilon)$
 $\vdash (\varepsilon, aB, 5)$
 $\vdash (\varepsilon, B, 54)$
 $\vdash (\varepsilon, S, 542)$




Example 9.6

$G :$

$S' \rightarrow S$	(1)
$S \rightarrow B \mid C$	(2, 3)
$B \rightarrow aB \mid b$	(4, 5)
$C \rightarrow aC \mid c$	(6, 7)

NBA(G):

$(ab, \varepsilon, \varepsilon)$
 $\vdash (b, a, \varepsilon)$
 $\vdash (\varepsilon, ab, \varepsilon)$
 $\vdash (\varepsilon, aB, 5)$
 $\vdash (\varepsilon, B, 54)$
 $\vdash (\varepsilon, S, 542)$
 $\vdash (\varepsilon, S', 5421)$



Definition 9.7 (LR(0) items and sets)

Let $G = \langle N, \Sigma, P, S \rangle \in CFG_{\Sigma}$ be start separated by $S' \rightarrow S$ and $S' \Rightarrow_r^* \alpha A w \Rightarrow_r \alpha \beta_1 \beta_2 w$ (i.e., $A \rightarrow \beta_1 \beta_2 \in P$).

- $[A \rightarrow \beta_1 \cdot \beta_2]$ is called an **LR(0) item** for $\alpha \beta_1$.

Definition 9.7 (LR(0) items and sets)

Let $G = \langle N, \Sigma, P, S \rangle \in CFG_\Sigma$ be start separated by $S' \rightarrow S$ and $S' \Rightarrow_r^* \alpha A w \Rightarrow_r \alpha \beta_1 \beta_2 w$ (i.e., $A \rightarrow \beta_1 \beta_2 \in P$).

- $[A \rightarrow \beta_1 \cdot \beta_2]$ is called an **LR(0) item** for $\alpha \beta_1$.
- Given $\gamma \in X^*$, $LR(0)(\gamma)$ denotes the set of all **LR(0)** items for γ , called the **LR(0) set** (or: **LR(0) information**) of γ .

Definition 9.7 (LR(0) items and sets)

Let $G = \langle N, \Sigma, P, S \rangle \in CFG_\Sigma$ be start separated by $S' \rightarrow S$ and $S' \Rightarrow_r^* \alpha A w \Rightarrow_r \alpha \beta_1 \beta_2 w$ (i.e., $A \rightarrow \beta_1 \beta_2 \in P$).

- $[A \rightarrow \beta_1 \cdot \beta_2]$ is called an **LR(0) item** for $\alpha \beta_1$.
- Given $\gamma \in X^*$, $LR(0)(\gamma)$ denotes the set of all **LR(0)** items for γ , called the **LR(0) set** (or: **LR(0) information**) of γ .
- $LR(0)(G) := \{LR(0)(\gamma) \mid \gamma \in X^*\}$.

Definition 9.7 (LR(0) items and sets)

Let $G = \langle N, \Sigma, P, S \rangle \in \text{CFG}_\Sigma$ be start separated by $S' \rightarrow S$ and $S' \Rightarrow_r^* \alpha A w \Rightarrow_r \alpha \beta_1 \beta_2 w$ (i.e., $A \rightarrow \beta_1 \beta_2 \in P$).

- $[A \rightarrow \beta_1 \cdot \beta_2]$ is called an **LR(0) item** for $\alpha \beta_1$.
- Given $\gamma \in X^*$, $\text{LR}(0)(\gamma)$ denotes the set of all **LR(0)** items for γ , called the **LR(0) set** (or: **LR(0) information**) of γ .
- $\text{LR}(0)(G) := \{\text{LR}(0)(\gamma) \mid \gamma \in X^*\}$.

Corollary 9.8

- 1 For every $\gamma \in X^*$, $\text{LR}(0)(\gamma)$ is finite.

Definition 9.7 (LR(0) items and sets)

Let $G = \langle N, \Sigma, P, S \rangle \in CFG_\Sigma$ be start separated by $S' \rightarrow S$ and $S' \Rightarrow_r^* \alpha A w \Rightarrow_r \alpha \beta_1 \beta_2 w$ (i.e., $A \rightarrow \beta_1 \beta_2 \in P$).

- $[A \rightarrow \beta_1 \cdot \beta_2]$ is called an **LR(0) item** for $\alpha \beta_1$.
- Given $\gamma \in X^*$, $LR(0)(\gamma)$ denotes the set of all **LR(0)** items for γ , called the **LR(0) set** (or: **LR(0) information**) of γ .
- $LR(0)(G) := \{LR(0)(\gamma) \mid \gamma \in X^*\}$.

Corollary 9.8

- 1 For every $\gamma \in X^*$, $LR(0)(\gamma)$ is finite.
- 2 $LR(0)(G)$ is finite.

Definition 9.7 (LR(0) items and sets)

Let $G = \langle N, \Sigma, P, S \rangle \in \text{CFG}_\Sigma$ be start separated by $S' \rightarrow S$ and $S' \Rightarrow_r^* \alpha A w \Rightarrow_r \alpha \beta_1 \beta_2 w$ (i.e., $A \rightarrow \beta_1 \beta_2 \in P$).

- $[A \rightarrow \beta_1 \cdot \beta_2]$ is called an **LR(0) item** for $\alpha \beta_1$.
- Given $\gamma \in X^*$, $\text{LR}(0)(\gamma)$ denotes the set of all **LR(0) items** for γ , called the **LR(0) set** (or: **LR(0) information**) of γ .
- $\text{LR}(0)(G) := \{\text{LR}(0)(\gamma) \mid \gamma \in X^*\}$.

Corollary 9.8

- 1 For every $\gamma \in X^*$, $\text{LR}(0)(\gamma)$ is finite.
- 2 $\text{LR}(0)(G)$ is finite.
- 3 The item $[A \rightarrow \beta \cdot] \in \text{LR}(0)(\gamma)$ indicates the possible **reduction** $(w, \alpha \beta, z) \vdash (w, \alpha A, zi)$ where $\pi_i = A \rightarrow \beta$ and $\gamma = \alpha \beta$.

Definition 9.7 (LR(0) items and sets)

Let $G = \langle N, \Sigma, P, S \rangle \in \text{CFG}_\Sigma$ be start separated by $S' \rightarrow S$ and $S' \Rightarrow_r^* \alpha A w \Rightarrow_r \alpha \beta_1 \beta_2 w$ (i.e., $A \rightarrow \beta_1 \beta_2 \in P$).

- $[A \rightarrow \beta_1 \cdot \beta_2]$ is called an **LR(0) item** for $\alpha \beta_1$.
- Given $\gamma \in X^*$, $\text{LR}(0)(\gamma)$ denotes the set of all **LR(0)** items for γ , called the **LR(0) set** (or: **LR(0) information**) of γ .
- $\text{LR}(0)(G) := \{\text{LR}(0)(\gamma) \mid \gamma \in X^*\}$.

Corollary 9.8

- 1 For every $\gamma \in X^*$, $\text{LR}(0)(\gamma)$ is finite.
- 2 $\text{LR}(0)(G)$ is finite.
- 3 The item $[A \rightarrow \beta \cdot] \in \text{LR}(0)(\gamma)$ indicates the possible **reduction** $(w, \alpha \beta, z) \vdash (w, \alpha A, z)$ where $\pi_i = A \rightarrow \beta$ and $\gamma = \alpha \beta$.
- 4 The item $[A \rightarrow \beta_1 \cdot Y \beta_2] \in \text{LR}(0)(\gamma)$ indicates an incomplete handle β_1 (to be completed by **shift** operations or ϵ -reductions).

Definition 9.9 (LR(0) conflicts)

Let $G = \langle N, \Sigma, P, S \rangle \in CFG_\Sigma$ and $I \in LR(0)(G)$.

- I has a **shift/reduce conflict** if there exist $A \rightarrow \alpha_1 a \alpha_2, B \rightarrow \beta \in P$ such that

$$[A \rightarrow \alpha_1 \cdot a \alpha_2], [B \rightarrow \beta \cdot] \in I.$$

Definition 9.9 ($LR(0)$ conflicts)

Let $G = \langle N, \Sigma, P, S \rangle \in CFG_{\Sigma}$ and $I \in LR(0)(G)$.

- I has a **shift/reduce conflict** if there exist $A \rightarrow \alpha_1 a \alpha_2, B \rightarrow \beta \in P$ such that

$$[A \rightarrow \alpha_1 \cdot a \alpha_2], [B \rightarrow \beta \cdot] \in I.$$

- I has a **reduce/reduce conflict** if there exist $A \rightarrow \alpha, B \rightarrow \beta \in P$ with $A \neq B$ or $\alpha \neq \beta$ such that

$$[A \rightarrow \alpha \cdot], [B \rightarrow \beta \cdot] \in I.$$

Definition 9.9 ($LR(0)$ conflicts)

Let $G = \langle N, \Sigma, P, S \rangle \in CFG_{\Sigma}$ and $I \in LR(0)(G)$.

- I has a **shift/reduce conflict** if there exist $A \rightarrow \alpha_1 a \alpha_2, B \rightarrow \beta \in P$ such that

$$[A \rightarrow \alpha_1 \cdot a \alpha_2], [B \rightarrow \beta \cdot] \in I.$$

- I has a **reduce/reduce conflict** if there exist $A \rightarrow \alpha, B \rightarrow \beta \in P$ with $A \neq B$ or $\alpha \neq \beta$ such that

$$[A \rightarrow \alpha \cdot], [B \rightarrow \beta \cdot] \in I.$$

Lemma 9.10

$G \in LR(0)$ iff no $I \in LR(0)(G)$ contains conflicting items.

Proof.

omitted □

Theorem 9.11 (Computing $LR(0)$ sets)

Let $G = \langle N, \Sigma, P, S \rangle \in CFG_{\Sigma}$ be start separated by $S' \rightarrow S$ and reduced.

- ① $LR(0)(\varepsilon)$ is the least set such that
- $[S' \rightarrow \cdot S] \in LR(0)(\varepsilon)$ and
 - if $[A \rightarrow \cdot B\gamma] \in LR(0)(\varepsilon)$ and $B \rightarrow \beta \in P$,
then $[B \rightarrow \cdot \beta] \in LR(0)(\varepsilon)$.

Theorem 9.11 (Computing $LR(0)$ sets)

Let $G = \langle N, \Sigma, P, S \rangle \in CFG_\Sigma$ be start separated by $S' \rightarrow S$ and reduced.

- ① $LR(0)(\varepsilon)$ is the least set such that
 - $[S' \rightarrow \cdot S] \in LR(0)(\varepsilon)$ and
 - if $[A \rightarrow \cdot B\gamma] \in LR(0)(\varepsilon)$ and $B \rightarrow \beta \in P$, then $[B \rightarrow \cdot \beta] \in LR(0)(\varepsilon)$.
- ② $LR(0)(\alpha Y)$ ($\alpha \in X^*, Y \in X$) is the least set such that
 - if $[A \rightarrow \gamma_1 \cdot Y\gamma_2] \in LR(0)(\alpha)$, then $[A \rightarrow \gamma_1 Y \cdot \gamma_2] \in LR(0)(\alpha Y)$ and
 - if $[A \rightarrow \gamma_1 \cdot B\gamma_2] \in LR(0)(\alpha Y)$ and $B \rightarrow \beta \in P$, then $[B \rightarrow \cdot \beta] \in LR(0)(\alpha Y)$.

Example 9.12 (cf. Example 9.6)

$G :$

S'	$\rightarrow$	S
S	$\rightarrow$	$B \mid C$
B	$\rightarrow$	$aB \mid b$
C	$\rightarrow$	$aC \mid c$

Example 9.12 (cf. Example 9.6)

$G :$

$S' \rightarrow S$	
$S \rightarrow B \mid C$	
$B \rightarrow aB \mid b$	$[S' \rightarrow \cdot S] \in LR(0)(\varepsilon)$
$C \rightarrow aC \mid c$	

$I_0 := LR(0)(\varepsilon) :$ $[S' \rightarrow \cdot S]$

Example 9.12 (cf. Example 9.6)

$G :$

$S' \rightarrow S$	
$S \rightarrow B \mid C$	$[A \rightarrow \cdot B\gamma] \in LR(0)(\varepsilon), B \rightarrow \beta \in P$
$B \rightarrow aB \mid b$	$\implies [B \rightarrow \cdot \beta] \in LR(0)(\varepsilon)$
$C \rightarrow aC \mid c$	

$I_0 := LR(0)(\varepsilon) :$

$[S' \rightarrow \cdot S]$	$[S \rightarrow \cdot B]$
----------------------------	---------------------------

Example 9.12 (cf. Example 9.6)

$G : \begin{array}{l} S' \rightarrow S \\ S \rightarrow B \mid C \\ B \rightarrow aB \mid b \\ C \rightarrow aC \mid c \end{array}$

$[A \rightarrow \cdot B \gamma] \in LR(0)(\varepsilon), B \rightarrow \beta \in P$

$\implies [B \rightarrow \cdot \beta] \in LR(0)(\varepsilon)$

$I_0 := LR(0)(\varepsilon) : \quad [S' \rightarrow \cdot S] \quad [S \rightarrow \cdot B] \quad [S \rightarrow \cdot C]$

Example 9.12 (cf. Example 9.6)

$G :$ $S' \rightarrow S$
 $S \rightarrow B \mid C$
 $B \rightarrow aB \mid b$
 $C \rightarrow aC \mid c$

$[A \rightarrow \cdot B\gamma] \in LR(0)(\varepsilon), B \rightarrow \beta \in P$
 $\implies [B \rightarrow \cdot \beta] \in LR(0)(\varepsilon)$

$I_0 := LR(0)(\varepsilon) :$ $\begin{bmatrix} S' \rightarrow \cdot S \\ B \rightarrow \cdot b \end{bmatrix}$ $[S \rightarrow \cdot B]$ $[S \rightarrow \cdot C]$ $[B \rightarrow \cdot aB]$

Example 9.12 (cf. Example 9.6)

$G :$ $S' \rightarrow S$
 $S \rightarrow B \mid C$
 $B \rightarrow aB \mid b$
 $C \rightarrow aC \mid c$

$[A \rightarrow \cdot B\gamma] \in LR(0)(\varepsilon), B \rightarrow \beta \in P$
 $\implies [B \rightarrow \cdot \beta] \in LR(0)(\varepsilon)$

$I_0 := LR(0)(\varepsilon) :$

$[S' \rightarrow \cdot S]$	$[S \rightarrow \cdot B]$	$[S \rightarrow \cdot C]$	$[B \rightarrow \cdot aB]$
$[B \rightarrow \cdot b]$	$[C \rightarrow \cdot aC]$	$[C \rightarrow \cdot c]$	

Example 9.12 (cf. Example 9.6)

$G : \begin{array}{l} S' \rightarrow S \\ S \rightarrow B \mid C \\ B \rightarrow aB \mid b \\ C \rightarrow aC \mid c \end{array}$

$$\begin{aligned} [A \rightarrow \gamma_1 \cdot Y \gamma_2] &\in LR(0)(\alpha) \\ \implies [A \rightarrow \gamma_1 Y \cdot \gamma_2] &\in LR(0)(\alpha Y) \end{aligned}$$

$$\begin{aligned} I_0 &:= LR(0)(\varepsilon) : & [S' \rightarrow \cdot S] & & [S \rightarrow \cdot B] & & [S \rightarrow \cdot C] & & [B \rightarrow \cdot aB] \\ & & [B \rightarrow \cdot b] & & [C \rightarrow \cdot aC] & & [C \rightarrow \cdot c] & & \\ I_1 &:= LR(0)(S) : & [S' \rightarrow S \cdot] & & & & & & \end{aligned}$$

Example 9.12 (cf. Example 9.6)

$G :$ $S' \rightarrow S$
 $S \rightarrow B \mid C$
 $B \rightarrow aB \mid b$
 $C \rightarrow aC \mid c$

$$\begin{aligned} [A \rightarrow \gamma_1 \cdot Y \gamma_2] &\in LR(0)(\alpha) \\ \implies [A \rightarrow \gamma_1 Y \cdot \gamma_2] &\in LR(0)(\alpha Y) \end{aligned}$$

$$\begin{aligned} I_0 &:= LR(0)(\varepsilon) : & [S' \rightarrow \cdot S] & & [S \rightarrow \cdot B] & & [S \rightarrow \cdot C] & & [B \rightarrow \cdot aB] \\ & & [B \rightarrow \cdot b] & & [C \rightarrow \cdot aC] & & [C \rightarrow \cdot c] & & \\ I_1 &:= LR(0)(S) : & [S' \rightarrow S \cdot] & & & & & & \\ I_2 &:= LR(0)(B) : & [S \rightarrow B \cdot] & & & & & & \end{aligned}$$

Example 9.12 (cf. Example 9.6)

$G :$ $S' \rightarrow S$
 $S \rightarrow B \mid C$
 $B \rightarrow aB \mid b$
 $C \rightarrow aC \mid c$

$$\begin{aligned} [A \rightarrow \gamma_1 \cdot Y \gamma_2] &\in LR(0)(\alpha) \\ \implies [A \rightarrow \gamma_1 Y \cdot \gamma_2] &\in LR(0)(\alpha Y) \end{aligned}$$

$$\begin{aligned} I_0 &:= LR(0)(\varepsilon) : & [S' \rightarrow \cdot S] & & [S \rightarrow \cdot B] & & [S \rightarrow \cdot C] & & [B \rightarrow \cdot aB] \\ & & [B \rightarrow \cdot b] & & [C \rightarrow \cdot aC] & & [C \rightarrow \cdot c] & & \\ I_1 &:= LR(0)(S) : & [S' \rightarrow S \cdot] & & & & & & \\ I_2 &:= LR(0)(B) : & [S \rightarrow B \cdot] & & & & & & \\ I_3 &:= LR(0)(C) : & [S \rightarrow C \cdot] & & & & & & \end{aligned}$$

Example 9.12 (cf. Example 9.6)

$G :$ $S' \rightarrow S$
 $S \rightarrow B \mid C$
 $B \rightarrow aB \mid b$
 $C \rightarrow aC \mid c$

$$\begin{aligned} [A \rightarrow \gamma_1 \cdot Y \gamma_2] &\in LR(0)(\alpha) \\ \implies [A \rightarrow \gamma_1 Y \cdot \gamma_2] &\in LR(0)(\alpha Y) \end{aligned}$$

$$\begin{aligned} I_0 &:= LR(0)(\varepsilon) : & [S' \rightarrow \cdot S] & & [S \rightarrow \cdot B] & & [S \rightarrow \cdot C] & & [B \rightarrow \cdot aB] \\ & & [B \rightarrow \cdot b] & & [C \rightarrow \cdot aC] & & [C \rightarrow \cdot c] & & \\ I_1 &:= LR(0)(S) : & [S' \rightarrow S \cdot] & & & & & & \\ I_2 &:= LR(0)(B) : & [S \rightarrow B \cdot] & & & & & & \\ I_3 &:= LR(0)(C) : & [S \rightarrow C \cdot] & & & & & & \\ I_4 &:= LR(0)(a) : & [B \rightarrow a \cdot B] & & [C \rightarrow a \cdot C] & & & & \end{aligned}$$

Example 9.12 (cf. Example 9.6)

$G :$ $S' \rightarrow S$
 $S \rightarrow B \mid C$
 $B \rightarrow aB \mid b$
 $C \rightarrow aC \mid c$

$[A \rightarrow \gamma_1 \cdot B \gamma_2] \in LR(0)(\alpha Y), B \rightarrow \beta \in P$
 $\implies [B \rightarrow \cdot \beta] \in LR(0)(\alpha Y)$

$I_0 := LR(0)(\varepsilon) :$ $[S' \rightarrow \cdot S]$ $[S \rightarrow \cdot B]$ $[S \rightarrow \cdot C]$ $[B \rightarrow \cdot aB]$
 $[B \rightarrow \cdot b]$ $[C \rightarrow \cdot aC]$ $[C \rightarrow \cdot c]$
 $I_1 := LR(0)(S) :$ $[S' \rightarrow S \cdot]$
 $I_2 := LR(0)(B) :$ $[S \rightarrow B \cdot]$
 $I_3 := LR(0)(C) :$ $[S \rightarrow C \cdot]$
 $I_4 := LR(0)(a) :$ $[B \rightarrow a \cdot B]$ $[C \rightarrow a \cdot C]$ $[B \rightarrow \cdot aB]$ $[B \rightarrow \cdot b]$

Example 9.12 (cf. Example 9.6)

$G : S' \rightarrow S$
 $S \rightarrow B \mid C$
 $B \rightarrow aB \mid b$
 $C \rightarrow aC \mid c$

$[A \rightarrow \gamma_1 \cdot B \gamma_2] \in LR(0)(\alpha Y), B \rightarrow \beta \in P$
 $\implies [B \rightarrow \cdot \beta] \in LR(0)(\alpha Y)$

$I_0 := LR(0)(\varepsilon) :$
 $\begin{bmatrix} S' \rightarrow \cdot S \\ B \rightarrow \cdot b \\ C \rightarrow \cdot aC \end{bmatrix}$
 $\begin{bmatrix} S \rightarrow \cdot B \\ C \rightarrow \cdot aC \end{bmatrix}$
 $\begin{bmatrix} S \rightarrow \cdot C \\ C \rightarrow \cdot c \end{bmatrix}$
 $[B \rightarrow \cdot aB]$

$I_1 := LR(0)(S) :$
 $[S' \rightarrow S \cdot]$

$I_2 := LR(0)(B) :$
 $[S \rightarrow B \cdot]$

$I_3 := LR(0)(C) :$
 $[S \rightarrow C \cdot]$

$I_4 := LR(0)(a) :$
 $\begin{bmatrix} B \rightarrow a \cdot B \\ C \rightarrow \cdot aC \end{bmatrix}$
 $\begin{bmatrix} C \rightarrow a \cdot C \\ C \rightarrow \cdot c \end{bmatrix}$
 $[B \rightarrow \cdot aB]$
 $[B \rightarrow \cdot b]$

Example 9.12 (cf. Example 9.6)

$G : S' \rightarrow S$
 $S \rightarrow B \mid C$
 $B \rightarrow aB \mid b$
 $C \rightarrow aC \mid c$

$$\begin{aligned} [A \rightarrow \gamma_1 \cdot Y \gamma_2] &\in LR(0)(\alpha) \\ \implies [A \rightarrow \gamma_1 Y \cdot \gamma_2] &\in LR(0)(\alpha Y) \end{aligned}$$

$I_0 := LR(0)(\varepsilon) :$ $[S' \rightarrow \cdot S]$ $[S \rightarrow \cdot B]$ $[S \rightarrow \cdot C]$ $[B \rightarrow \cdot aB]$
 $[B \rightarrow \cdot b]$ $[C \rightarrow \cdot aC]$ $[C \rightarrow \cdot c]$

$I_1 := LR(0)(S) :$ $[S' \rightarrow S \cdot]$

$I_2 := LR(0)(B) :$ $[S \rightarrow B \cdot]$

$I_3 := LR(0)(C) :$ $[S \rightarrow C \cdot]$

$I_4 := LR(0)(a) :$ $[B \rightarrow a \cdot B]$ $[C \rightarrow a \cdot C]$ $[B \rightarrow \cdot aB]$ $[B \rightarrow \cdot b]$
 $[C \rightarrow \cdot aC]$ $[C \rightarrow \cdot c]$

$I_5 := LR(0)(b) :$ $[B \rightarrow b \cdot]$

Example 9.12 (cf. Example 9.6)

$G : S' \rightarrow S$
 $S \rightarrow B \mid C$
 $B \rightarrow aB \mid b$
 $C \rightarrow aC \mid c$

$$\begin{aligned} [A \rightarrow \gamma_1 \cdot Y \gamma_2] &\in LR(0)(\alpha) \\ \implies [A \rightarrow \gamma_1 Y \cdot \gamma_2] &\in LR(0)(\alpha Y) \end{aligned}$$

$I_0 := LR(0)(\varepsilon) :$ $\begin{bmatrix} S' \rightarrow \cdot S \\ B \rightarrow \cdot b \\ C \rightarrow \cdot aC \end{bmatrix}$ $\begin{bmatrix} S \rightarrow \cdot B \\ C \rightarrow \cdot aC \end{bmatrix}$ $\begin{bmatrix} S \rightarrow \cdot C \\ C \rightarrow \cdot c \end{bmatrix}$ $[B \rightarrow \cdot aB]$

$I_1 := LR(0)(S) :$ $\begin{bmatrix} S' \rightarrow S \cdot \end{bmatrix}$

$I_2 := LR(0)(B) :$ $\begin{bmatrix} S \rightarrow B \cdot \end{bmatrix}$

$I_3 := LR(0)(C) :$ $\begin{bmatrix} S \rightarrow C \cdot \end{bmatrix}$

$I_4 := LR(0)(a) :$ $\begin{bmatrix} B \rightarrow a \cdot B \\ C \rightarrow a \cdot C \end{bmatrix}$ $\begin{bmatrix} C \rightarrow a \cdot C \\ C \rightarrow \cdot c \end{bmatrix}$ $[B \rightarrow \cdot aB]$ $[B \rightarrow \cdot b]$

$I_5 := LR(0)(b) :$ $\begin{bmatrix} B \rightarrow b \cdot \end{bmatrix}$

$I_6 := LR(0)(c) :$ $\begin{bmatrix} C \rightarrow c \cdot \end{bmatrix}$

Example 9.12 (cf. Example 9.6)

$G : S' \rightarrow S$
 $S \rightarrow B \mid C$
 $B \rightarrow aB \mid b$
 $C \rightarrow aC \mid c$

$$\begin{aligned}
 &[A \rightarrow \gamma_1 \cdot Y \gamma_2] \in LR(0)(\alpha) \\
 \implies &[A \rightarrow \gamma_1 Y \cdot \gamma_2] \in LR(0)(\alpha Y)
 \end{aligned}$$

$I_0 := LR(0)(\varepsilon) :$
 $\begin{array}{llll} [S' \rightarrow \cdot S] & [S \rightarrow \cdot B] & [S \rightarrow \cdot C] & [B \rightarrow \cdot aB] \\ [B \rightarrow \cdot b] & [C \rightarrow \cdot aC] & [C \rightarrow \cdot c] & \end{array}$

$I_1 := LR(0)(S) :$
 $[S' \rightarrow S \cdot]$

$I_2 := LR(0)(B) :$
 $[S \rightarrow B \cdot]$

$I_3 := LR(0)(C) :$
 $[S \rightarrow C \cdot]$

$I_4 := LR(0)(a) :$
 $\begin{array}{llll} [B \rightarrow a \cdot B] & [C \rightarrow a \cdot C] & [B \rightarrow \cdot aB] & [B \rightarrow \cdot b] \\ [C \rightarrow \cdot aC] & [C \rightarrow \cdot c] & \end{array}$

$I_5 := LR(0)(b) :$
 $[B \rightarrow b \cdot]$

$I_6 := LR(0)(c) :$
 $[C \rightarrow c \cdot]$

$I_7 := LR(0)(aB) :$
 $[B \rightarrow aB \cdot]$

Example 9.12 (cf. Example 9.6)

$G : S' \rightarrow S$
 $S \rightarrow B \mid C$
 $B \rightarrow aB \mid b$
 $C \rightarrow aC \mid c$

$$\begin{aligned}
 &[A \rightarrow \gamma_1 \cdot Y \gamma_2] \in LR(0)(\alpha) \\
 \implies &[A \rightarrow \gamma_1 Y \cdot \gamma_2] \in LR(0)(\alpha Y)
 \end{aligned}$$

$I_0 := LR(0)(\varepsilon) :$
 $\begin{array}{llll} [S' \rightarrow \cdot S] & [S \rightarrow \cdot B] & [S \rightarrow \cdot C] & [B \rightarrow \cdot aB] \\ [B \rightarrow \cdot b] & [C \rightarrow \cdot aC] & [C \rightarrow \cdot c] & \end{array}$

$I_1 := LR(0)(S) :$
 $[S' \rightarrow S \cdot]$

$I_2 := LR(0)(B) :$
 $[S \rightarrow B \cdot]$

$I_3 := LR(0)(C) :$
 $[S \rightarrow C \cdot]$

$I_4 := LR(0)(a) :$
 $\begin{array}{llll} [B \rightarrow a \cdot B] & [C \rightarrow a \cdot C] & [B \rightarrow \cdot aB] & [B \rightarrow \cdot b] \\ [C \rightarrow \cdot aC] & [C \rightarrow \cdot c] & \end{array}$

$I_5 := LR(0)(b) :$
 $[B \rightarrow b \cdot]$

$I_6 := LR(0)(c) :$
 $[C \rightarrow c \cdot]$

$I_7 := LR(0)(aB) :$
 $[B \rightarrow aB \cdot]$

$I_8 := LR(0)(aC) :$
 $[C \rightarrow aC \cdot]$

Example 9.12 (cf. Example 9.6)

$G : S' \rightarrow S$
 $S \rightarrow B \mid C$
 $B \rightarrow aB \mid b$
 $C \rightarrow aC \mid c$

$I_0 := LR(0)(\varepsilon) :$ $[S' \rightarrow \cdot S]$ $[S \rightarrow \cdot B]$ $[S \rightarrow \cdot C]$ $[B \rightarrow \cdot aB]$
 $[B \rightarrow \cdot b]$ $[C \rightarrow \cdot aC]$ $[C \rightarrow \cdot c]$

$I_1 := LR(0)(S) :$ $[S' \rightarrow S \cdot]$

$I_2 := LR(0)(B) :$ $[S \rightarrow B \cdot]$

$I_3 := LR(0)(C) :$ $[S \rightarrow C \cdot]$

$I_4 := LR(0)(a) :$ $[B \rightarrow a \cdot B]$ $[C \rightarrow a \cdot C]$ $[B \rightarrow \cdot aB]$ $[B \rightarrow \cdot b]$
 $[C \rightarrow \cdot aC]$ $[C \rightarrow \cdot c]$

$I_5 := LR(0)(b) :$ $[B \rightarrow b \cdot]$

$I_6 := LR(0)(c) :$ $[C \rightarrow c \cdot]$

$I_7 := LR(0)(aB) :$ $[B \rightarrow aB \cdot]$

$I_8 := LR(0)(aC) :$ $[C \rightarrow aC \cdot]$

$(LR(0)(aa) = LR(0)(a) = I_4, LR(0)(ab) = LR(0)(b) = I_5,$
 $LR(0)(ac) = LR(0)(c) = I_6, I_9 := LR(0)(\gamma) = \emptyset$ in all remaining cases)

Example 9.12 (cf. Example 9.6)

$$\begin{aligned} G : \quad & S' \rightarrow S \\ & S \rightarrow B \mid C \\ & B \rightarrow aB \mid b \\ & C \rightarrow aC \mid c \end{aligned}$$
$$I_0 := LR(0)(\varepsilon) : \quad \begin{array}{llll} [S' \rightarrow \cdot S] & [S \rightarrow \cdot B] & [S \rightarrow \cdot C] & [B \rightarrow \cdot aB] \\ [B \rightarrow \cdot b] & [C \rightarrow \cdot aC] & [C \rightarrow \cdot c] & \end{array}$$
$$I_1 := LR(0)(S) : \quad [S' \rightarrow S \cdot]$$
$$I_2 := LR(0)(B) : \quad [S \rightarrow B \cdot]$$
$$I_3 := LR(0)(C) : \quad [S \rightarrow C \cdot]$$
$$I_4 := LR(0)(a) : \quad \begin{array}{llll} [B \rightarrow a \cdot B] & [C \rightarrow a \cdot C] & [B \rightarrow \cdot aB] & [B \rightarrow \cdot b] \\ [C \rightarrow \cdot aC] & [C \rightarrow \cdot c] & & \end{array}$$
$$I_5 := LR(0)(b) : \quad [B \rightarrow b \cdot]$$
$$I_6 := LR(0)(c) : \quad [C \rightarrow c \cdot]$$
$$I_7 := LR(0)(aB) : \quad [B \rightarrow aB \cdot]$$
$$I_8 := LR(0)(aC) : \quad [C \rightarrow aC \cdot]$$

$(LR(0)(aa) = LR(0)(a) = I_4, LR(0)(ab) = LR(0)(b) = I_5,$
 $LR(0)(ac) = LR(0)(c) = I_6, I_9 := LR(0)(\gamma) = \emptyset$ in all remaining cases)

no conflicts $\implies G \in LR(0)$

- 1 Repetition: Nondeterministic Bottom-Up Parsing
- 2 Nondeterministic Bottom-Up Parsing (cont.)
- 3 $LR(k)$ Grammars
- 4 $LR(0)$ Grammars
- 5 Examples of $LR(0)$ Conflicts

Example 9.13

$G :$ $S' \rightarrow S$
 $S \rightarrow Aa \mid Bb$
 $A \rightarrow a$
 $B \rightarrow a$

Reduce/Reduce Conflicts

Example 9.13

$G : S' \rightarrow S$
 $S \rightarrow Aa \mid Bb$
 $A \rightarrow a$
 $B \rightarrow a$

$LR(0)(\varepsilon) : [S' \rightarrow \cdot S] \quad [S \rightarrow \cdot Aa] \quad [S \rightarrow \cdot Bb] \quad [A \rightarrow \cdot a] \quad [B \rightarrow \cdot a]$
 $LR(0)(S) : [S' \rightarrow S \cdot]$
 $LR(0)(A) : [S \rightarrow A \cdot a]$
 $LR(0)(B) : [S \rightarrow B \cdot a]$
 $LR(0)(a) : [A \rightarrow a \cdot] \quad [B \rightarrow a \cdot]$
 $LR(0)(Aa) : [S \rightarrow Aa \cdot]$
 $LR(0)(Ba) : [S \rightarrow Ba \cdot]$

Example 9.13

$G : S' \rightarrow S$
 $S \rightarrow Aa \mid Bb$
 $A \rightarrow a$
 $B \rightarrow a$

$LR(0)(\varepsilon) : [S' \rightarrow \cdot S] \quad [S \rightarrow \cdot Aa] \quad [S \rightarrow \cdot Bb] \quad [A \rightarrow \cdot a] \quad [B \rightarrow \cdot a]$
 $LR(0)(S) : [S' \rightarrow S \cdot]$
 $LR(0)(A) : [S \rightarrow A \cdot a]$
 $LR(0)(B) : [S \rightarrow B \cdot a]$
 $LR(0)(a) : [A \rightarrow a \cdot] \quad [B \rightarrow a \cdot]$
 $LR(0)(Aa) : [S \rightarrow Aa \cdot]$
 $LR(0)(Ba) : [S \rightarrow Ba \cdot]$

Note: G is unambiguous

Example 9.14

$G : S' \rightarrow S$
 $S \rightarrow aS \mid a$

Example 9.14

$G : S' \rightarrow S$
 $S \rightarrow aS \mid a$

$LR(0)(\varepsilon) : [S' \rightarrow \cdot S] \quad [S \rightarrow \cdot aS] \quad [S \rightarrow \cdot a]$

$LR(0)(S) : [S' \rightarrow S \cdot]$

$LR(0)(a) : [S \rightarrow a \cdot S] \quad [S \rightarrow \cdot aS] \quad [S \rightarrow \cdot a] \quad [S \rightarrow a \cdot]$

$LR(0)(aS) : [S \rightarrow aS \cdot]$

Example 9.14

$G : S' \rightarrow S$
 $S \rightarrow aS \mid a$

$LR(0)(\varepsilon) : [S' \rightarrow \cdot S] \quad [S \rightarrow \cdot aS] \quad [S \rightarrow \cdot a]$

$LR(0)(S) : [S' \rightarrow S \cdot]$

$LR(0)(a) : [S \rightarrow a \cdot S] \quad [S \rightarrow \cdot aS] \quad [S \rightarrow \cdot a] \quad [S \rightarrow a \cdot]$

$LR(0)(aS) : [S \rightarrow aS \cdot]$

Note: G is unambiguous