

Semantics and Verification of Software

Lecture 2: Operational Semantics of WHILE I (Evaluation of Expressions)

Thomas Noll

Lehrstuhl für Informatik 2
(Software Modeling and Verification)

RWTH Aachen University

`noll@cs.rwth-aachen.de`

`http://www-i2.informatik.rwth-aachen.de/i2/svsw10/`

Summer Semester 2010

- 1 Repetition: Syntax of WHILE
- 2 Operational Semantics of WHILE
- 3 Evaluation of Arithmetic Expressions
- 4 Excursus: Proof by Structural Induction
- 5 Evaluation of Boolean Expressions

WHILE: simple imperative programming language without procedures or advanced data structures

Syntactic categories:

Category	Domain	Meta variable
Numbers	$\mathbb{Z} = \{0, 1, -1, \dots\}$	z
Truth values	$\mathbb{B} = \{\text{true}, \text{false}\}$	t
Variables	$Var = \{x, y, \dots\}$	x
Arithmetic expressions	$AExp$ (next slide)	a
Boolean expressions	$BExp$ (next slide)	b
Commands (statements)	Cmd (next slide)	c

Syntax of WHILE Programs

Definition (Syntax of WHILE)

The **syntax of WHILE Programs** is defined by the following context-free grammar:

$$a ::= z \mid x \mid a_1 + a_2 \mid a_1 - a_2 \mid a_1 * a_2 \in AExp$$
$$b ::= t \mid a_1 = a_2 \mid a_1 > a_2 \mid \neg b \mid b_1 \wedge b_2 \mid b_1 \vee b_2 \in BExp$$
$$c ::= \text{skip} \mid x := a \mid c_1 ; c_2 \mid \text{if } b \text{ then } c_1 \text{ else } c_2 \mid \text{while } b \text{ do } c \in Cmd$$

Remarks: we assume that

- the syntax of numbers, truth values and variables is predefined (i.e., no “lexical analysis”)
- the syntax of ambiguous constructs is uniquely determined (by brackets, priorities, or indentation)

- 1 Repetition: Syntax of WHILE
- 2 Operational Semantics of WHILE
- 3 Evaluation of Arithmetic Expressions
- 4 Excursus: Proof by Structural Induction
- 5 Evaluation of Boolean Expressions

Operational Semantics of WHILE

- Idea: define meaning of programs by specifying its behaviour being executed on an (abstract) machine

Operational Semantics of WHILE

- Idea: define meaning of programs by specifying its behaviour being executed on an (abstract) machine
- Here: evaluation/execution relation for program fragments (expressions, statements)

Operational Semantics of WHILE

- Idea: define meaning of programs by specifying its behaviour being executed on an (abstract) machine
- Here: evaluation/execution relation for program fragments (expressions, statements)
- Approach based on Structural Operational Semantics (SOS)

G.D. Plotkin: A structural approach to operational semantics, DAIMI FN-19, Computer Science Department, Aarhus University, 1981

Operational Semantics of WHILE

- Idea: define meaning of programs by specifying its behaviour being executed on an (abstract) machine
- Here: evaluation/execution relation for program fragments (expressions, statements)
- Approach based on Structural Operational Semantics (SOS)

G.D. Plotkin: A structural approach to operational semantics, DAIMI FN-19, Computer Science Department, Aarhus University, 1981

- Employs derivation rules of the form

$$\text{Name} \frac{\text{Premise(s)}}{\text{Conclusion}}$$

- meaning: if every premise is fulfilled, then conclusion can be drawn
- a rule with no premises is called an axiom

Operational Semantics of WHILE

- Idea: define meaning of programs by specifying its behaviour being executed on an (abstract) machine
- Here: evaluation/execution relation for program fragments (expressions, statements)
- Approach based on Structural Operational Semantics (SOS)

G.D. Plotkin: A structural approach to operational semantics, DAIMI FN-19, Computer Science Department, Aarhus University, 1981

- Employs derivation rules of the form

$$\text{Name} \frac{\text{Premise(s)}}{\text{Conclusion}}$$

- meaning: if every premise is fulfilled, then conclusion can be drawn
 - a rule with no premises is called an axiom
- Derivation rules can be composed to form derivation trees with axioms as leafs (formal definition later)

- 1 Repetition: Syntax of WHILE
- 2 Operational Semantics of WHILE
- 3 Evaluation of Arithmetic Expressions
- 4 Excursus: Proof by Structural Induction
- 5 Evaluation of Boolean Expressions

- Meaning of expression = value (in the usual sense)
- Depends on the values of the variables in the expression

- **Meaning of expression** = value (in the usual sense)
- Depends on the **values of the variables** in the expression

Definition 2.1 (Program state)

A **(program) state** is an element of the set

$$\Sigma := \{\sigma \mid \sigma : Var \rightarrow \mathbb{Z}\},$$

called the **state space**.

Thus $\sigma(x)$ denotes the value of $x \in Var$ in state $\sigma \in \Sigma$.

Evaluation of Arithmetic Expressions I

Remember: $a ::= z \mid x \mid a_1 + a_2 \mid a_1 - a_2 \mid a_1 * a_2 \in AExp$

Evaluation of Arithmetic Expressions I

Remember: $a ::= z \mid x \mid a_1 + a_2 \mid a_1 - a_2 \mid a_1 * a_2 \in AExp$

Definition 2.2 (Evaluation relation for arithmetic expressions)

If $a \in AExp$ and $\sigma \in \Sigma$, then $\langle a, \sigma \rangle$ is called a **configuration**.

Expression a **evaluates to** $z \in \mathbb{Z}$ in state σ (notation: $\langle a, \sigma \rangle \rightarrow z$) if this relationship is derivable by means of the following rules:

Axioms:

$$\frac{}{\langle z, \sigma \rangle \rightarrow z} \quad \frac{}{\langle x, \sigma \rangle \rightarrow \sigma(x)}$$

Rules:

$$\frac{\langle a_1, \sigma \rangle \rightarrow z_1 \quad \langle a_2, \sigma \rangle \rightarrow z_2}{\langle a_1 + a_2, \sigma \rangle \rightarrow z} \quad \text{where } z := z_1 + z_2$$

$$\frac{\langle a_1, \sigma \rangle \rightarrow z_1 \quad \langle a_2, \sigma \rangle \rightarrow z_2}{\langle a_1 - a_2, \sigma \rangle \rightarrow z} \quad \text{where } z := z_1 - z_2$$

$$\frac{\langle a_1, \sigma \rangle \rightarrow z_1 \quad \langle a_2, \sigma \rangle \rightarrow z_2}{\langle a_1 * a_2, \sigma \rangle \rightarrow z} \quad \text{where } z := z_1 * z_2$$

Example 2.3

$a = (x+3)*(y-2)$, $\sigma(x) = 3$, $\sigma(y) = 9$:

Example 2.3

$a = (x+3)*(y-2)$, $\sigma(x) = 3$, $\sigma(y) = 9$:

$$\frac{}{\langle (x+3)*(y-2), \sigma \rangle \rightarrow}$$

Example 2.3

$a = (x+3)*(y-2)$, $\sigma(x) = 3$, $\sigma(y) = 9$:

$$\frac{\frac{}{\langle x+3, \sigma \rangle \rightarrow} \quad \frac{}{\langle y-2, \sigma \rangle \rightarrow}}{\langle (x+3)*(y-2), \sigma \rangle \rightarrow}$$

$$\frac{\langle a_1, \sigma \rangle \rightarrow z_1 \quad \langle a_2, \sigma \rangle \rightarrow z_2}{\langle a_1 * a_2, \sigma \rangle \rightarrow z} \quad \text{where } z := z_1 * z_2$$

Example 2.3

$a = (x+3)*(y-2)$, $\sigma(x) = 3$, $\sigma(y) = 9$:

$$\frac{\frac{\overline{\langle x, \sigma \rangle \rightarrow} \quad \overline{\langle 3, \sigma \rangle \rightarrow}}{\overline{\langle x+3, \sigma \rangle \rightarrow}} \quad \overline{\langle y-2, \sigma \rangle \rightarrow}}{\overline{\langle (x+3)*(y-2), \sigma \rangle \rightarrow}}$$

$$\frac{\overline{\langle a_1, \sigma \rangle \rightarrow z_1} \quad \overline{\langle a_2, \sigma \rangle \rightarrow z_2}}{\overline{\langle a_1+a_2, \sigma \rangle \rightarrow z}} \quad \text{where } z := z_1 + z_2$$

Example 2.3

$a = (x+3)*(y-2)$, $\sigma(\mathbf{x}) = 3$, $\sigma(y) = 9$:

$$\begin{array}{c}
 \frac{\langle x, \sigma \rangle \rightarrow 3 \quad \langle 3, \sigma \rangle \rightarrow}{\langle x+3, \sigma \rangle \rightarrow} \quad \frac{}{\langle y-2, \sigma \rangle \rightarrow} \\
 \hline
 \langle (x+3)*(y-2), \sigma \rangle \rightarrow \\
 \\
 \hline
 \langle x, \sigma \rangle \rightarrow \sigma(x)
 \end{array}$$

Example 2.3

$a = (x+3)*(y-2)$, $\sigma(x) = 3$, $\sigma(y) = 9$:

$$\frac{\frac{\overline{\langle x, \sigma \rangle \rightarrow 3} \quad \overline{\langle 3, \sigma \rangle \rightarrow 3}}{\overline{\langle x+3, \sigma \rangle \rightarrow}} \quad \frac{\overline{\quad} \quad \overline{\quad}}{\overline{\langle y-2, \sigma \rangle \rightarrow}}}{\overline{\langle (x+3)*(y-2), \sigma \rangle \rightarrow}}$$
$$\frac{\overline{\quad}}{\overline{\langle z, \sigma \rangle \rightarrow z}}$$

Example 2.3

$a = (x+3)*(y-2)$, $\sigma(x) = 3$, $\sigma(y) = 9$:

$$\frac{\frac{\overline{\langle x, \sigma \rangle \rightarrow 3} \quad \overline{\langle 3, \sigma \rangle \rightarrow 3}}{\overline{\langle x+3, \sigma \rangle \rightarrow 6}} \quad \frac{\overline{\quad} \quad \overline{\quad}}{\overline{\langle y-2, \sigma \rangle \rightarrow \quad}}}{\overline{\langle (x+3)*(y-2), \sigma \rangle \rightarrow \quad}}$$

$$\frac{\overline{\langle a_1, \sigma \rangle \rightarrow z_1} \quad \overline{\langle a_2, \sigma \rangle \rightarrow z_2}}{\overline{\langle a_1+a_2, \sigma \rangle \rightarrow z}} \quad \text{where } z := z_1 + z_2$$

Example 2.3

$a = (x+3)*(y-2)$, $\sigma(x) = 3$, $\sigma(y) = 9$:

$$\frac{\frac{\overline{\langle x, \sigma \rangle \rightarrow 3} \quad \overline{\langle 3, \sigma \rangle \rightarrow 3}}{\overline{\langle x+3, \sigma \rangle \rightarrow 6}} \quad \frac{\overline{\langle y, \sigma \rangle \rightarrow} \quad \overline{\langle 2, \sigma \rangle \rightarrow}}{\overline{\langle y-2, \sigma \rangle \rightarrow}}}{\overline{\langle (x+3)*(y-2), \sigma \rangle \rightarrow}}$$

$$\frac{\overline{\langle a_1, \sigma \rangle \rightarrow z_1} \quad \overline{\langle a_2, \sigma \rangle \rightarrow z_2}}{\overline{\langle a_1 - a_2, \sigma \rangle \rightarrow z}} \quad \text{where } z := z_1 - z_2$$

Example 2.3

$a = (x+3)*(y-2)$, $\sigma(x) = 3$, $\sigma(y) = 9$:

$$\frac{\frac{\overline{\langle x, \sigma \rangle \rightarrow 3} \quad \overline{\langle 3, \sigma \rangle \rightarrow 3}}{\overline{\langle x+3, \sigma \rangle \rightarrow 6}} \quad \frac{\overline{\langle y, \sigma \rangle \rightarrow 9} \quad \overline{\langle 2, \sigma \rangle \rightarrow}}{\overline{\langle y-2, \sigma \rangle \rightarrow}}}{\overline{\langle (x+3)*(y-2), \sigma \rangle \rightarrow}}$$
$$\overline{\langle x, \sigma \rangle \rightarrow \sigma(x)}$$

Example 2.3

$a = (x+3)*(y-2)$, $\sigma(x) = 3$, $\sigma(y) = 9$:

$$\frac{\frac{\overline{\langle x, \sigma \rangle \rightarrow 3} \quad \overline{\langle 3, \sigma \rangle \rightarrow 3}}{\langle x+3, \sigma \rangle \rightarrow 6} \quad \frac{\overline{\langle y, \sigma \rangle \rightarrow 9} \quad \overline{\langle 2, \sigma \rangle \rightarrow 2}}{\langle y-2, \sigma \rangle \rightarrow 7}}{\langle (x+3)*(y-2), \sigma \rangle \rightarrow 42}$$
$$\frac{}{\langle z, \sigma \rangle \rightarrow z}$$

Example 2.3

$a = (x+3)*(y-2)$, $\sigma(x) = 3$, $\sigma(y) = 9$:

$$\frac{\frac{\overline{\langle x, \sigma \rangle \rightarrow 3} \quad \overline{\langle 3, \sigma \rangle \rightarrow 3}}{\overline{\langle x+3, \sigma \rangle \rightarrow 6}} \quad \frac{\overline{\langle y, \sigma \rangle \rightarrow 9} \quad \overline{\langle 2, \sigma \rangle \rightarrow 2}}{\overline{\langle y-2, \sigma \rangle \rightarrow 7}}}{\overline{\langle (x+3)*(y-2), \sigma \rangle \rightarrow}}$$

$$\frac{\overline{\langle a_1, \sigma \rangle \rightarrow z_1} \quad \overline{\langle a_2, \sigma \rangle \rightarrow z_2}}{\overline{\langle a_1 - a_2, \sigma \rangle \rightarrow z}} \quad \text{where } z := z_1 - z_2$$

Example 2.3

$a = (x+3)*(y-2)$, $\sigma(x) = 3$, $\sigma(y) = 9$:

$$\frac{\frac{\overline{\langle x, \sigma \rangle \rightarrow 3} \quad \overline{\langle 3, \sigma \rangle \rightarrow 3}}{\langle x+3, \sigma \rangle \rightarrow 6} \quad \frac{\overline{\langle y, \sigma \rangle \rightarrow 9} \quad \overline{\langle 2, \sigma \rangle \rightarrow 2}}{\langle y-2, \sigma \rangle \rightarrow 7}}{\langle (x+3)*(y-2), \sigma \rangle \rightarrow 42}$$

$$\frac{\langle a_1, \sigma \rangle \rightarrow z_1 \quad \langle a_2, \sigma \rangle \rightarrow z_2}{\langle a_1 * a_2, \sigma \rangle \rightarrow z} \quad \text{where } z := z_1 * z_2$$

Example 2.3

$a = (x+3)*(y-2)$, $\sigma(x) = 3$, $\sigma(y) = 9$:

$$\frac{\frac{\overline{\langle x, \sigma \rangle \rightarrow 3} \quad \overline{\langle 3, \sigma \rangle \rightarrow 3}}{\overline{\langle x+3, \sigma \rangle \rightarrow 6}} \quad \frac{\overline{\langle y, \sigma \rangle \rightarrow 9} \quad \overline{\langle 2, \sigma \rangle \rightarrow 2}}{\overline{\langle y-2, \sigma \rangle \rightarrow 7}}}{\overline{\langle (x+3)*(y-2), \sigma \rangle \rightarrow 42}}$$

Here: structure of derivation tree = structure of program fragment
(generally not true)

First formal result: value of an expression only depends on valuation of variables which occur (freely) in the expression

First formal result: value of an expression only depends on valuation of variables which occur (freely) in the expression

Definition 2.4 (Free variables)

The set of **free variables** of an expression is given by the function

$$FV : AExp \rightarrow 2^{Var}$$

where

$$\begin{array}{ll} FV(z) := \emptyset & FV(a_1 + a_2) := FV(a_1) \cup FV(a_2) \\ FV(x) := \{x\} & FV(a_1 - a_2) := FV(a_1) \cup FV(a_2) \\ & FV(a_1 * a_2) := FV(a_1) \cup FV(a_2) \end{array}$$

Free Variables

First formal result: value of an expression only depends on valuation of variables which occur (freely) in the expression

Definition 2.4 (Free variables)

The set of **free variables** of an expression is given by the function

$$FV : AExp \rightarrow 2^{Var}$$

where

$$\begin{array}{ll} FV(z) := \emptyset & FV(a_1 + a_2) := FV(a_1) \cup FV(a_2) \\ FV(x) := \{x\} & FV(a_1 - a_2) := FV(a_1) \cup FV(a_2) \\ & FV(a_1 * a_2) := FV(a_1) \cup FV(a_2) \end{array}$$

Result will be shown by **structural induction** on the expression

- 1 Repetition: Syntax of WHILE
- 2 Operational Semantics of WHILE
- 3 Evaluation of Arithmetic Expressions
- 4 Excursus: Proof by Structural Induction
- 5 Evaluation of Boolean Expressions

Proof principle

Given: an inductive set, i.e., a set S

- which contains certain atomic elements and
- which is closed under certain operations

To show: property $P(s)$ applies to every $s \in S$

Proof: we verify:

Induction base: $P(s)$ holds for every atomic element s

Induction hypothesis: assume that $P(s_1), P(s_2)$ etc.

Induction step: then also $P(f(s_1, \dots, s_n))$ holds for every operation f of arity n

Application: natural numbers (“complete induction”)

Definition: $\mathbb{N}$ is the least set which

- contains 0 and
- contains $n + 1$ whenever $n \in \mathbb{N}$

Induction base: $P(0)$ holds

Induction hypothesis: $P(n)$ holds

Induction step: $P(n + 1)$ holds

Example 2.5

We prove that $P(n) : \sum_{i=1}^n i = \frac{n(n+1)}{2}$ holds for every $n \in \mathbb{N}$.

Example 2.5

We prove that $P(n) : \sum_{i=1}^n i = \frac{n(n+1)}{2}$ holds for every $n \in \mathbb{N}$.

$P(0)$ holds: $\sum_{i=1}^0 i = 0 = \frac{0(0+1)}{2} \quad \checkmark$

Example 2.5

We prove that $P(n) : \sum_{i=1}^n i = \frac{n(n+1)}{2}$ holds for every $n \in \mathbb{N}$.

$P(0)$ holds: $\sum_{i=1}^0 i = 0 = \frac{0(0+1)}{2} \quad \checkmark$

Assume $P(n)$: $\sum_{i=1}^n i = \frac{n(n+1)}{2}$

Example 2.5

We prove that $P(n) : \sum_{i=1}^n i = \frac{n(n+1)}{2}$ holds for every $n \in \mathbb{N}$.

$P(0)$ holds: $\sum_{i=1}^0 i = 0 = \frac{0(0+1)}{2} \quad \checkmark$

Assume $P(n)$: $\sum_{i=1}^n i = \frac{n(n+1)}{2}$

Show $P(n+1)$: $\sum_{i=1}^{n+1} i = \sum_{i=1}^n i + (n+1)$

Example 2.5

We prove that $P(n) : \sum_{i=1}^n i = \frac{n(n+1)}{2}$ holds for every $n \in \mathbb{N}$.

$P(0)$ holds: $\sum_{i=1}^0 i = 0 = \frac{0(0+1)}{2} \quad \checkmark$

Assume $P(n)$: $\sum_{i=1}^n i = \frac{n(n+1)}{2}$

Show $P(n+1)$: $\sum_{i=1}^{n+1} i = \sum_{i=1}^n i + (n+1)$
 $= \frac{n(n+1)}{2} + (n+1)$

Example 2.5

We prove that $P(n) : \sum_{i=1}^n i = \frac{n(n+1)}{2}$ holds for every $n \in \mathbb{N}$.

$P(0)$ holds: $\sum_{i=1}^0 i = 0 = \frac{0(0+1)}{2} \quad \checkmark$

Assume $P(n)$: $\sum_{i=1}^n i = \frac{n(n+1)}{2}$

Show $P(n+1)$: $\sum_{i=1}^{n+1} i = \sum_{i=1}^n i + (n+1)$
 $= \frac{n(n+1)}{2} + (n+1)$
 $= \frac{n(n+1)}{2} + \frac{2(n+1)}{2}$

Example 2.5

We prove that $P(n) : \sum_{i=1}^n i = \frac{n(n+1)}{2}$ holds for every $n \in \mathbb{N}$.

$P(0)$ holds: $\sum_{i=1}^0 i = 0 = \frac{0(0+1)}{2} \quad \checkmark$

Assume $P(n)$: $\sum_{i=1}^n i = \frac{n(n+1)}{2}$

Show $P(n+1)$: $\sum_{i=1}^{n+1} i = \sum_{i=1}^n i + (n+1)$
 $= \frac{n(n+1)}{2} + (n+1)$
 $= \frac{n(n+1)}{2} + \frac{2(n+1)}{2}$
 $= \frac{(n+2)(n+1)}{2}$

Example 2.5

We prove that $P(n) : \sum_{i=1}^n i = \frac{n(n+1)}{2}$ holds for every $n \in \mathbb{N}$.

$P(0)$ holds: $\sum_{i=1}^0 i = 0 = \frac{0(0+1)}{2} \quad \checkmark$

Assume $P(n)$: $\sum_{i=1}^n i = \frac{n(n+1)}{2}$

Show $P(n+1)$: $\sum_{i=1}^{n+1} i = \sum_{i=1}^n i + (n+1)$

$$\begin{aligned} &= \frac{n(n+1)}{2} + (n+1) \\ &= \frac{n(n+1)}{2} + \frac{2(n+1)}{2} \\ &= \frac{(n+2)(n+1)}{2} \\ &= \frac{(n+1)((n+1)+1)}{2} \quad \checkmark \end{aligned}$$

Application: arithmetic expressions (Def. 1.2)

Definition: $AExp$ is the least set which

- contains all integers $z \in \mathbb{Z}$ and all variables $x \in Var$ and
- contains $a_1 + a_2$, $a_1 - a_2$ and $a_1 * a_2$ whenever $a_1, a_2 \in AExp$

Induction base: $P(z)$ and $P(x)$ holds (for every $z \in \mathbb{Z}$ and $x \in Var$)

Induction hypothesis: $P(a_1)$ and $P(a_2)$ holds

Induction step: $P(a_1 + a_2)$, $P(a_1 - a_2)$ and $P(a_1 * a_2)$ holds

Lemma 2.6

Let $a \in AExp$ and $\sigma, \sigma' \in \Sigma$ such that $\sigma(x) = \sigma'(x)$ for every $x \in FV(a)$. Then, for every $z \in \mathbb{Z}$,

$$\langle a, \sigma \rangle \rightarrow z \iff \langle a, \sigma' \rangle \rightarrow z.$$

Lemma 2.6

Let $a \in AExp$ and $\sigma, \sigma' \in \Sigma$ such that $\sigma(x) = \sigma'(x)$ for every $x \in FV(a)$. Then, for every $z \in \mathbb{Z}$,

$$\langle a, \sigma \rangle \rightarrow z \iff \langle a, \sigma' \rangle \rightarrow z.$$

Proof.

by **structural induction** on a (on the board)



- 1 Repetition: Syntax of WHILE
- 2 Operational Semantics of WHILE
- 3 Evaluation of Arithmetic Expressions
- 4 Excursus: Proof by Structural Induction
- 5 Evaluation of Boolean Expressions

Evaluation of Boolean Expressions I

Remember: $b ::= t \mid a_1 = a_2 \mid a_1 > a_2 \mid \neg b \mid b_1 \wedge b_2 \mid b_1 \vee b_2 \in BExp$

Evaluation of Boolean Expressions I

Remember: $b ::= t \mid a_1 = a_2 \mid a_1 > a_2 \mid \neg b \mid b_1 \wedge b_2 \mid b_1 \vee b_2 \in BExp$

Definition 2.7 (Evaluation relation for Boolean expressions)

For $b \in BExp$, $\sigma \in \Sigma$, and $t \in \mathbb{B}$, the **evaluation relation** $\langle b, \sigma \rangle \rightarrow t$ is defined by the following rules:

$$\overline{\langle t, \sigma \rangle \rightarrow t}$$

$$\frac{\langle a_1, \sigma \rangle \rightarrow z \quad \langle a_2, \sigma \rangle \rightarrow z}{\langle a_1 = a_2, \sigma \rangle \rightarrow \text{true}}$$

$$\langle a_1 = a_2, \sigma \rangle \rightarrow \text{true}$$

$$\frac{\langle a_1, \sigma \rangle \rightarrow z_1 \quad \langle a_2, \sigma \rangle \rightarrow z_2}{\langle a_1 > a_2, \sigma \rangle \rightarrow \text{true}} \text{ if } z_1 > z_2$$

$$\langle a_1 > a_2, \sigma \rangle \rightarrow \text{true}$$

$$\frac{\langle b, \sigma \rangle \rightarrow \text{false}}{\langle \neg b, \sigma \rangle \rightarrow \text{true}}$$

$$\langle \neg b, \sigma \rangle \rightarrow \text{true}$$

$$\frac{\langle b_1, \sigma \rangle \rightarrow \text{true} \quad \langle b_2, \sigma \rangle \rightarrow \text{true}}{\langle b_1 \wedge b_2, \sigma \rangle \rightarrow \text{true}}$$

$$\langle b_1 \wedge b_2, \sigma \rangle \rightarrow \text{true}$$

$$\frac{\langle b_1, \sigma \rangle \rightarrow \text{false} \quad \langle b_2, \sigma \rangle \rightarrow \text{true}}{\langle b_1 \wedge b_2, \sigma \rangle \rightarrow \text{false}}$$

$$\langle b_1 \wedge b_2, \sigma \rangle \rightarrow \text{false}$$

$$\frac{\langle a_1, \sigma \rangle \rightarrow z_1 \quad \langle a_2, \sigma \rangle \rightarrow z_2}{\langle a_1 = a_2, \sigma \rangle \rightarrow \text{false}} \text{ if } z_1 \neq z_2$$

$$\langle a_1 = a_2, \sigma \rangle \rightarrow \text{false}$$

$$\frac{\langle a_1, \sigma \rangle \rightarrow z_1 \quad \langle a_2, \sigma \rangle \rightarrow z_2}{\langle a_1 > a_2, \sigma \rangle \rightarrow \text{false}} \text{ if } z_1 \leq z_2$$

$$\langle a_1 > a_2, \sigma \rangle \rightarrow \text{false}$$

$$\frac{\langle b, \sigma \rangle \rightarrow \text{true}}{\langle \neg b, \sigma \rangle \rightarrow \text{false}}$$

$$\langle \neg b, \sigma \rangle \rightarrow \text{false}$$

$$\frac{\langle b_1, \sigma \rangle \rightarrow \text{true} \quad \langle b_2, \sigma \rangle \rightarrow \text{false}}{\langle b_1 \wedge b_2, \sigma \rangle \rightarrow \text{false}}$$

$$\langle b_1 \wedge b_2, \sigma \rangle \rightarrow \text{false}$$

$$\frac{\langle b_1, \sigma \rangle \rightarrow \text{false} \quad \langle b_2, \sigma \rangle \rightarrow \text{false}}{\langle b_1 \wedge b_2, \sigma \rangle \rightarrow \text{false}}$$

$$\langle b_1 \wedge b_2, \sigma \rangle \rightarrow \text{false}$$

($\vee$ analogously)

Remarks:

- Binary Boolean operators $\wedge$ and $\vee$ are interpreted as **strict**, i.e., always evaluate both arguments.

Important in situations like

```
while p <> nil and p^.key < val do ...!
```

(see next slides for alternatives)

Remarks:

- Binary Boolean operators $\wedge$ and $\vee$ are interpreted as **strict**, i.e., always evaluate both arguments.

Important in situations like

```
while p <> nil and p^.key < val do ...!
```

(see next slides for alternatives)

- $FV : BExp \rightarrow 2^{Var}$ can be defined in analogy to Def. 2.4.

Remarks:

- Binary Boolean operators $\wedge$ and $\vee$ are interpreted as **strict**, i.e., always evaluate both arguments.

Important in situations like

```
while p <> nil and p^.key < val do ...!
```

(see next slides for alternatives)

- $FV : BExp \rightarrow 2^{Var}$ can be defined in analogy to Def. 2.4.
- Lemma 2.6 holds analogously for Boolean expressions, i.e., the value of $b \in BExp$ does not depend on variables in $Var \setminus FV(b)$.

Definition 2.8 (Sequential evaluation of Boolean expressions)

For $b \in BExp$, $\sigma \in \Sigma$, and $t \in \mathbb{B}$, the **sequential evaluation relation** $\langle b, \sigma \rangle \rightarrow t$ is defined by the following rules:

$$\begin{array}{c} \vdots \\ \frac{\langle b_1, \sigma \rangle \rightarrow \text{false}}{\langle b_1 \wedge b_2, \sigma \rangle \rightarrow \text{false}} \quad \frac{\langle b_1, \sigma \rangle \rightarrow \text{true} \quad \langle b_2, \sigma \rangle \rightarrow t}{\langle b_1 \wedge b_2, \sigma \rangle \rightarrow t} \\ \frac{\langle b_1, \sigma \rangle \rightarrow \text{true}}{\langle b_1 \vee b_2, \sigma \rangle \rightarrow \text{true}} \quad \frac{\langle b_1, \sigma \rangle \rightarrow \text{false} \quad \langle b_2, \sigma \rangle \rightarrow t}{\langle b_1 \vee b_2, \sigma \rangle \rightarrow t} \end{array}$$

Evaluation of Boolean Expressions IV

Definition 2.9 (Parallel evaluation of Boolean expressions)

For $b \in BExp$, $\sigma \in \Sigma$, and $t \in \mathbb{B}$, the **parallel evaluation relation** $\langle b, \sigma \rangle \rightarrow t$ is defined by the following rules:

$\vdots$

$$\frac{\langle b_1, \sigma \rangle \rightarrow \text{false}}{\langle b_1 \wedge b_2, \sigma \rangle \rightarrow \text{false}} \quad \frac{\langle b_2, \sigma \rangle \rightarrow \text{false}}{\langle b_1 \wedge b_2, \sigma \rangle \rightarrow \text{false}}$$

$$\frac{\langle b_1, \sigma \rangle \rightarrow \text{true} \quad \langle b_2, \sigma \rangle \rightarrow \text{true}}{\langle b_1 \wedge b_2, \sigma \rangle \rightarrow \text{true}}$$

$$\frac{\langle b_1, \sigma \rangle \rightarrow \text{true}}{\langle b_1 \vee b_2, \sigma \rangle \rightarrow \text{true}} \quad \frac{\langle b_2, \sigma \rangle \rightarrow \text{true}}{\langle b_1 \vee b_2, \sigma \rangle \rightarrow \text{true}}$$

$$\frac{\langle b_1, \sigma \rangle \rightarrow \text{false} \quad \langle b_2, \sigma \rangle \rightarrow \text{false}}{\langle b_1 \vee b_2, \sigma \rangle \rightarrow \text{false}}$$